



Spectroscopic Investigation Complex Formation Of Vanadium Using 2,6-Dithiol-4-Methylphenol And Hydrophob Amins

Kerim A. Kuliyev

Department of Chemistry, Azerbaijan State Pedagogical University, U. Gadhibekov street 34. Baku, AZ 1000
Azerbaijan

kerim.kuliev.69@mail.ru

ABSTRACT

Mixed-ligand complexes of Vanadium (IV, V) with 2,6-dithiol-4-methylphenol (DTMP) and hydrofob amins (Am) have been studied by spectrophotometry. Extraction of mixed ligand complexes is maximal at pH 3.8-5.3. The optimal conditions for the formation and extraction of mixed-ligand compounds have been found and the ratios of components in the complexes have been determined. The Beer's law was applicable in the range of 0.04-3.6 g/ml. The method is free from common interferences. A procedure has been developed for extraction – spectrophotometric determination vanadium in soils and in oil and oil-products.

Indexing terms/Keywords

Extraction-spectrophotometric method, vanadium, 2,6-dithiol-4-methylphenol (DTMP), hydrofob amins (Am), mixed-ligand complexes.

Academic Discipline And Sub-Disciplines

Chemistry

SUBJECT CLASSIFICATION

Analytical chemistry; Inorganic Chemistry; General chemistry

TYPE (METHOD/APPROACH)

Types of research: experiments; Approach: quantitative research

Council for Innovative Research

Peer Review Research Publishing System

Journal: Journal of Advances in Chemistry

Vol. 11, No. 4

editorjaonline@gmail.com

www.cirjac.com



1. Introduction

Vanadium is a trace element and is of great industrial, ecological, biological and pharmacological importance. Vanadium metal is present in different alloys and is an important element present in steels. In view of crucial role in different fields, it is necessary to develop sensitive analytical procedures for the determination of trace amounts of vanadium.

Survey of literature reveals that several procedures have been reported for the determination of vanadium. The very sensitive, fairly selective direct spectrophotometric method for the determination of trace amount of V(V) with naphthalene-1,5-diamine has been developed. The method was successfully applied in the determination of vanadium in environmental, and biological samples [1].

Novel, rapid, highly sensitive and selective spectrophotometric method for the determination of traces of V(V) in environmental and biological samples, pharmaceutical and steel samples was studied. The method is based on oxidation of 2,4-dinitro phenyl hydrazine (2,4-DNPH) by V(V) followed by coupling reaction with N-(1-naphthalene-1-yl)ethane-1,2-diamine-dihydrochloride (NEDA) in acidic medium to give red colored derivative or oxidation of 4-Amino Pyridine by V(V) followed by coupling reaction with NEDA in basic medium to give pink colored derivative [2]. Several kinetic methods based on its catalytic action on the oxidation of organic compounds have been reported in literature [3-5]. The bromates have been extensively used as the oxidizing agents [4, 5], although hydrogen peroxide and potassium chlorate have also been used for the same purpose [6]. In the present paper, a simple and sensitive method is proposed for V(IV) determination in the presence of V(V). This is based on the oxidation of V(IV) present in the sample to V(V) by addition of Fe(III) cation, followed by a complexation reaction of Fe(III) with the spectrophotometric reagent 2-(5-bromo-2-pyridylazo)-5-diethylaminophenol (Br-PADAP) [7].

In scrutiny of literature reveals that several spectrophotometric methods have been reported for the determination of vanadium in environmental and biological samples. Recently, few authors introduced various reagents for spectrophotometric determination of Vanadium in various samples such as 2-(2-quinolyazo)-5-diethylaminophenol [8], variamine blue [9], cryamine R [10], Benzylacetate [11], pyrogallol [12], 2-(8-quinolyazo)-5-dimethylamino-phenol [13], 2-hydroxyacetophenone [14], 4-(2-pyridylazo) resorcinol [15], tannic acid [16], 2-(5-chloro-2-pyridylazo)-5-dimethylaminophenol [17] and *N,N*-bis(2-hydroxy-3-sulfo-propyl)-toluidine [18].

Oxyphenolate complexes of Vanadium are insoluble in chloroform, while mixed-ligand complexes with hydrophobic amines and aminophenols easily dissolve in various organic solvents [19, 20, 21]. In this respect, a very promising reagent is 2,6-dithiol-4-methylphenol (DTMP), which contains one hydroxyl and two sulfhydryl groups and is a sulfur-containing analogue of mononuclear polyphenols with two oxygen atoms replaced with sulfur atoms.

The present study is concerned with the investigation of V(IV, V) interaction with 2,6-dithiol-4-methylphenol (DTMP), resulting in the formation of colored complexes insoluble in nonpolar solvents. Experiments on electromigration in a U-shaped tube and on sorption on EDE-10P (EDE- ethylenediamine, epichlorohydrin; 10- serial number of the brand: P- means that the matrix has a macroporous structure) anion exchangers have demonstrated the anionic nature of single-ligand complexes, in the electromigration study of the complexes, it was found that the blue-green dithiophenolate complexes of Vanadium (IV, V) moved to the cathode. When the sign of the charge of the single-ligand complexes was determined by ion chromatography, the EDE-10P anion exchanger completely absorbed the colored component of the solution. When hydrophobic amines (Am) were introduced into the system, the extraction of these compounds into the organic phase as a mixed-ligand complex (MLC) was observed. Among hydrophobic amines, *o*-toluidine (*o*-tol), *m*-toluidine (*m*-tol) and *p*-toluidine (*p*-tol) were used. Based on these data, new selective and highly sensitive procedures were developed for the extraction-spectrophotometric determination of trace vanadium in soils and in oil and oil-products.

2. Experimental

2.1. Instruments

The absorbance of the extracts was measured using a KFK-2 photocolormeter and an SF-26 spectrophotometer; the equilibrium value of the pH of aqueous phase was measured using a I-120.2 potentiometer with a glass electrode. ESR spectra of solutions of mixed-ligand complexes were registered on a JEOS-JES-PE-3X spectrometer (Japan) with working a frequency of 9400 MHz. IR - spectra were recorded on a spectrophotometer UR-20.

2.2. Reagents and solutions.

Stock solutions (1.96×10^{-2} M) of V(IV, V) were prepared from chemically pure salts $\text{VOSO}_4 \cdot 3\text{H}_2\text{O}$ and $\text{NaVO}_3 \cdot 2\text{H}_2\text{O}$. The working solutions with concentration 0.1 mg/mL were obtained by the dilution of the stock solutions. The concentration of solutions of Vanadium(V) and vanadium(IV) was determined by titration with iron(II) salts and potassium permanganate, respectively [22]. Solutions of DTMP and Am in chloroform (0.01 M) were used. To create the optimal acidity, 0.1 M solutions of KOH and HCl were used. The extractant was purified chloroform.

2.3. General procedure

2.3.1. General procedure for the determination of Vanadium (IV, V).

Portions of stock solutions of Vanadium (IV, V) varying from 0.1 to 1.0 mL with a 0.1-mL step, a 2.0-mL portion of a 0.01 M solution of DTMP, and a 2.0 mL portion of a 0.01 M solution of Am were placed in to calibrated test tubes with ground-glass stoppers (the volume of the organic phase was 5 mL). The volume of the organic phase was increased to 5



mL using chloroform. The required value of pH was adjusted by adding 0.1M HCl. The volume of the aqueous phase was increased to 20 mL using distilled water. In 10 min after the complete separation of the phases, the organic phase was separated from the aqueous phase and the absorbance of the extracts was measured on KFK-2 at room temperature and 590 nm ($l=0.5\text{cm}$).

2.3.2. Determination of Vanadium in soils

The proposed procedures for the determination of Vanadium were applied to its determination in light-chestnut soil from the Caspian zone. A 0.5-g weight was finely ground in an agate mortar and calcined in muffle furnace for 3 h. after cooling, the sample was treated and dissolved in an graphite cup in a mixture of 16 mL of conc. HF, 5mL of conc. HNO_3 , and 15 mL of conc. HCl at 50-60°C. to remove excess hydrogen fluoride, a 8 mL portion of conc. HNO_3 was added triply to the solution that was each time evaporated to 5-6mL. After that, the solution was transferred into a 100 mL volumetric flask and its volume was brought to the mark with distilled water. Vanadium was determined in aliquot portions of the solution using the proposed procedures.

2.3.3. Determination of Vanadium in oil and oil-products of Baku

For this purpose, we took a china cup (9 cm diameter) of used fuel (40g) and burned it. Twenty grams of fuel oil or 10 grams of hydron ($\pm 0.0001\text{g}$) were used. The China cup was heated at $500\pm 20^\circ\text{C}$ for one hour. After cooling 5 mL of HCl(1:1), 2 mL of fuel, and 0,5g Na_2CO_3 without water was added to the cup and warmed. The cup was heated for 2 to 3 minutes in an oven at 800°C . After cooling 5 mL of distilled water was added to the mass in the cup. It was poured twice through the same filter into a 50-mL graduated cylinder to a volume of 15mL. We then added a little vanadyl-sulphate with 10 mg of vanadium, 2-3 mL of concentrate HCl and corrected the volume with distilled water.

To a given quantity of this solution, we added 0.5 g of thiocarbamide, 2 mL of 0.01 M DTMP and 2.2 mL of 0.01 M Am. The volume of the organic phase was increased to 5 mL using chloroform. After shaking we increased the volume of solution to 25 mL with distilled water. We separated the organic-phase from the water-phase and sured the optical absorbency of the extract at 590 nm using the CFC-2. The concentration of Vanadium was determined and corrected statistically.

3. Results and discussion

In weakly acid and neutral media (pH 1.4-7.5) V (IV,V) forms with DTMP a blue compound soluble in water. When the concentration of the reagent and pH were increased, the color became more intense. Being organic cations, hydrophobamins under certain conditions in acid media form intensely colored MLC with the acido complexes of V (IV,V). The formed complexes are extractable with organic solvents.

3.1. The choice of the extractant

For the extraction of complexes we used chloroform, tetrachloromethane, benzene, toluene, xylene, isobutanol, isopentanol, and dichloroethane. The extractivity of the complexes was estimated by the distribution coefficient and recovery. Chloroform, dichloroethane, and chloro-benzene appeared to be the best extractants. All the further investigations were carried out with chloroform. The concentration of vanadium in the organic phase was determined with 8-hydroxyquinoline [23] by photometric measurements after back extraction, while in the aqueous phase it was determined by the difference. The basicity of AP hardly influences the recovery of vanadium. After a single extraction with chloroform, 98.4-98.9% of vanadium was extracted as an ion associate.

3.2. Influence of the pH of the aqueous phase

The effect of pH on the intensity of the color reaction is shown in the Fig. 1. The absorbance was found to be maximum in the pH range 3,8-5,3. Hence further analytical investigations were carried out in media of pH 4. Extraction of V(IV,V) enhanced with the increase in the acidity of the initial solution; the further increase in acidity lead to the gradual decrease of recovery, which was obviously associated with a decrease in the concentration of the ionized form of DTMP. Probably, it is present in the solution in the non-dissociated state. At $\text{pH}\geq 7$, the complexes were hardly extracted, obviously because of the decrease in the degree of Am protonation (Fig.1).




Fig.1. Absorbance of mixed-ligand complexes as a function of the pH of the aqueous phase.

1. V(IV)-DTMP-*o*-tol; 2. V(IV)-DTMP-*m*-tol; 3. V(IV)-DTBP-*p*-tol.

$C_V=3.92 \times 10^{-5}$ M, $C_{DTMP}=0.8 \times 10^{-3}$ M, $C_{Am}=0.88 \times 10^{-3}$ M, KFK-2, 590nm, $l=0.5$ cm.

3.3. Influence of reagent concentration and incubation time

For the formation and extraction of MLC, a 20-25-fold excess of complexing reagents is required; for example, the optimal conditions for formation and extraction of these compounds are provided by 0.8×10^{-3} M DTMP and 0.88×10^{-3} M Am. A large excess of hydrophob amin interferers with the determination. However it was found that the presence of excess of the reagent solution does not alter the absorbance of the color reaction.

Unlike single-ligand complexes, mixed-ligand complexes of V (IV, V) with DTMP and Am were stable in aqueous and organic solvents and did not decompose for two days, or over a month after extraction. The required duration of the phase contact was 10 min. the system obeyed the Beer's law in the range 0.04-3.6 mg/mL of vanadium.

3.4. Electronic absorption spectra

Neither the metal ion nor the reagent has appreciable absorbance at specified wavelengths. Hence further studies were carried out at 620-630 nm (fig.2). The molar absorption coefficient was $(3.2-3.7) \times 10^4$ L · mol⁻¹ cm⁻¹.




Fig.2. Absorption of mixed-ligand complexes

1. V(IV)-DTMP-*o*-tol; 2. V(IV)-DTMP-*m*-tol; 3. V(IV)-DTBP-*p*-tol.

$C_V=3.92 \times 10^{-5}$ M, $C_{DTMP}=0.8 \times 10^{-3}$ M, $C_{Am}=0.88 \times 10^{-3}$ M, SF-26, $l=1$ cm.

3.5. Stoichiometry of the complexes and the mechanism of complexation

It was found using the Nazarenko method that V(IV) in the complexes was present in the form of VO^{2+} . The number of protons replaced by vanadium in one DTBP molecule appeared to be two [24,25].

It was found that the spectrophotometric characteristics of the MLC of V(IV) and V(V) were identical, i.e., in the interaction with DTMP, V(V) was reduced to V(IV) and VO^{2+} was the complex-producing form. This fact was also confirmed by ESR spectrometry [26]. V(V) does not have unpaired electrons and is diamagnetic, while V(IV) has one d electron and exhibits electron paramagnetic absorption. According to the value of the nuclear spin $j=7/2$, the ESR spectra of V(IV) consisted of eight lines with the hyperfine structure associated with the interaction of the magnetic moment of an unpaired electron with the magnetic moment of the ^{51}V nucleus. Hyperfine structure consisting of 8 lines was observed in the ESR spectra of chloroform extracts of MLC from aqueous solutions of V(IV) and V(V) salts. Consequently, the interaction of V(IV) and V(V) with DTMP involved its reduction to V(IV). The results of the studies are presented in fig.3.

Spectrophotometric data and ESR spectra allow us to present the mechanism of interaction in the studied systems. V(V) was firstly reduced by 2,6-dithiolphenol to V(IV), the later then participated in the complexation reaction. Therefore, at the optimal pH, DTMP formed a colored complex of anionic nature with V(IV) and V(V), which was transferred to the organic phase as a mixed-ligand complex with a quadratic pyramide structure as a result of the addition of Am.




Fig.3. Hyperfine splitting of the paramagnetic resonance line in solutions:

(1) V(IV)-DTMP-o-tol and (2) V(V)-DTMP-o-tol.

The stoichiometric coefficients of the reaction were determined by the Asmus method and by the equilibrium shift method [27]. The formation of MLC can be presented in the following way. When vanadium ion interact with two molecules of DTMP, they form doubly-charged anionic complexes, which are extracted with two molecules of protonated Am (Fig.4).

Constant of stability of complexes are determined by method of crossing of curves [27].




Fig.4. Determination of the ratio of components by the equilibrium shift method for (a) V(IV)-DTMP-o-tol and (b) V(IV)-DTMP-m-tol. 1- V:DTMP; 2- V : Am.

$C_V = 3.92 \times 10^{-5} \text{ M}$. SF-26, $l = 590 \text{ nm}$, $l = 1 \text{ cm}$

The IR spectra of the complexes in the field of $790\text{--}810 \text{ cm}^{-1}$ there is an intensive strip of absorption caused by valent vibration of group $[V = O]^{2+}$. The disappearance of a distinct strip at 2580 cm^{-1} , observed in ranges DTMP and emergence spectrum of complexes of two strips of absorption one of which is displaced towards smaller frequencies, says that one of the sulfhydryl groups participates in formation of a complex. Observed reduction of intensity of a strip of absorption in area $3200\text{--}3600 \text{ cm}^{-1}$ with a maximum at 3450 cm^{-1} , emergence of a wide strip in area $3000\text{--}3100 \text{ cm}^{-1}$ shows that the hydroxyl group takes part in formation of coordination communication in the ionized state. Detection of strips of absorption at 1380 cm^{-1} indicates availability of the protonated tolidine [28,29]. Structure extractable complexes can be represented as in Figure 5.




Fig. 5. Structure of complex

3.6. Influence of interfering ions

Large amounts of alkali and alkaline-earth metals and REE do not interfere with the determination of vanadium. The interference of Fe(III) was eliminated by thioglycolic acid or 20% solution of SnCl_2 ; Cu(II), Cr(VI), and Mn(VII) were masked by thiourea; Ti(IV), ascorbic acid; and Zr(IV), Nb(V), and Ta(V), by fluoride ions. Ions of Mo(VI), W(VI), Ti(IV), Nb(V), and Ta(V) form with DTMP and Am colored compounds and interfere with the determination of vanadium. However, these elements form complexes in more acidic medium. Since Mo(VI) and W(VI) complexes with DTMP are soluble in chloroform, while complexes of V(IV) are insoluble, the latter ones are completely transferred into the aqueous



phase as V-DTMP. After the separation of the aqueous phase from the organic phase, vanadium was determined in the aqueous phase under optimal conditions with the addition of Am. The selectivity of the photometric determination of V(IV) with DTMP is demonstrated in Table 1. In conclusion the analytical parameters pertaining to the proposed method are given in the Table 2.

Table 1. Influence of interfering ions on the determination of V(IV) as MLC with DTMP and *p*-toluidine (30,0 mg V added)




Ion	Molar excess of the ion	Masking agent	Found V, μg	RSD, %
Co(II)	150		30,0	3
Ni(II)	150		30,2	3
Al(III)	190		29,8	4
Fe(II)	300		29,6	3
Fe(III)	50	Thioglycolic acid (HS-CS-COOH)	30,2	4
Cd(II)	200		30,2	3
Zr(IV)	75	NaF	30,5	5
Cu(II)	30	SC(NH ₂) ₂	29,7	5
Ti(IV)	25	Ascorbic acid (C ₆ H ₈ O ₆)	29,6	4
W(VI)	15		29,8	4
Mo(VI)	5		29,6	5
Cr(III)	30		30,2	2
Nb(V)	25	NaF	30,2	5
Ta(V)	25	NaF	30,2	5
Pt(II)	50		29,2	5
Pd(II)	50		30,4	6
UO ₂ ²⁺	45	CH ₃ COO ⁻	29,3	4

Table 2. Optical characteristics, precision and accuracy of the spectrophotometric determination of V(IV) with DTMP and toluidines

Parameter	<i>o</i> -toluidine (<i>o</i> -tol)	<i>m</i> -toluidine (<i>m</i> -tol)	<i>p</i> -toluidine (<i>p</i> -tol)
Color	blue-green	blue-green	blue-green
The pH range of education and extraction	3,4-5,8	3,5-6,0	3,7-6,1
The pH range of maximum extraction	3,8-4,7	3,9-5,1	4,0-5,3
λ_{max} (nm)	620	625	630
Molar absorptivity (L · mol ⁻¹ cm ⁻¹)	3.2 · 10 ⁴	3.4 · 10 ⁴	3.7 · 10 ⁴
Sandell's sensitivity ($\mu\text{g} \cdot \text{cm}^{-2}$)	0.016	0.015	0.014
R, %	98.4	98.6	98.9
The equation of calibration curves	0.032+0.61x	0.035+0.65x	0.032+0.71x
Correlation coefficient	0.9985	0.9982	0.9976
lg k _e	5.85	5.92	6.00
Stability constant (β)	9.1	9.5	9.8
Beer's law range ($\mu\text{g} \cdot \text{ml}^{-1}$)	0.05-3.2	0.05-3.4	0.04-3.6

Table 3 demonstrates the data which allow a comparison of the analytical parameters of the procedures for the determination of Vanadium with the earlier known procedures [23,30].

Table 3. Comparative characteristics of the procedures for determining vanadium

Reagent	pH (solvent)	λ , nm	$\epsilon \cdot 10^{-4}$	Beer's law range
 Sulfonitrazo	2,3	522	2,06	
 8- Mercaptoquinoline	4,0 - 5,5(chloroform, toluene)	412	0,30	0,5-1,5
 8 - Hydroxyquinoline	3,0 – 5,0 (chloroform)	550	3,00	-
DTMP+ <i>o</i> -tol	3.8-4.7(chloroform)	620	3.2	0.05-3.2
DTMP+ <i>m</i> -tol	3.9-5.1(chloroform)	625	3,4	0,05-3.4

3.7. Effect of vanadium (IV) concentration

The adherence to Beer's law was studied by measuring the absorbance value of the series of solutions containing different concentrations of the metal ion. A linear calibration graph drawn between absorbance and the metal ion concentration indicates that V(V) may be determined in the range 0.04-3.6 $\mu\text{g/ml}$. The pertaining calibration graph is shown in the Fig. 6.




Fig. 6. Analytical determination of V (IV)

$C_V=3.92 \times 10^{-5}$ M, $C_{DTMP}=1.0 \times 10^{-3}$ M, $C_{Am}=0.8 \times 10^{-3}$ M, KFK-2, 590nm, $l=0.5$ cm.

3.8. Analytical Applications

The proposed method under the already established optimum conditions was applied for the determination of V(V) in soils and in oil and oil-products. The proposed procedures for determining Vanadium in soils were verified by 8-hydroxyquinoline and formaldoxime methods [23]. The results of the analysis are listed in Table 4 indicate the successful applicability of the proposed method to real sample analysis.

Table 4. Determination of Vanadium in various samples (n=6, P=0.95)

Prosedure	\bar{X} , %	RSD, %	$\bar{X} \pm \frac{t_p \cdot S}{\sqrt{n}}$
<i>Soil</i>			
Standard method			
8 - Hydroxyquinoline	$1,13 \cdot 10^{-2}$	2.0	$(1,13 \pm 0.024) \cdot 10^{-2}$
H-C=N-OH Formaldoksim	$1,14 \cdot 10^{-2}$	2.3	$(1,14 \pm 0.027) \cdot 10^{-2}$
Proposed method			
DTMP+o-tol	$1,13 \cdot 10^{-2}$	2.2	$(1,13 \pm 0.024) \cdot 10^{-2}$
DTMP+m-tol	$1,12 \cdot 10^{-2}$	1.6	$(1,12 \pm 0.019) \cdot 10^{-2}$
DTMP+p-tol	$1,11 \cdot 10^{-2}$	1.8	$(1,11 \pm 0.021) \cdot 10^{-2}$
<i>Oil</i>			
Standard method			



8 - Hydroxyquinoline	$6.18 \cdot 10^{-5}$	1.8	$(6.18 \pm 0.12) \cdot 10^{-5}$
Proposed method			
DTMP+o-tol	$6.12 \cdot 10^{-5}$	1.5	$(6.12 \pm 0.09) \cdot 10^{-5}$
<i>Fuel Oil</i>			
Standard method			
Formaldoksim	$2.58 \cdot 10^{-3}$	1.6	$(2.58 \pm 0.04) \cdot 10^{-3}$
Proposed method			
DTMP+m-tol	$2.56 \cdot 10^{-3}$	1.3	$(2.56 \pm 0.03) \cdot 10^{-3}$
<i>Hydron</i>			
Standard method			
8 - Hydroxyquinoline	$4.31 \cdot 10^{-3}$	1.5	$(4.31 \pm 0.07) \cdot 10^{-3}$
Proposed method			
DTMP+m-tol	$4.28 \cdot 10^{-3}$	1.3	$(4.28 \pm 0.06) \cdot 10^{-3}$
DTMP+p-tol	$4.27 \cdot 10^{-3}$	1.4	$(4.27 \pm 0.06) \cdot 10^{-3}$

CONCLUSIONS

Mixed-ligand complexes of Vanadium (IV, V) with 2,6-dithiol-4-methylphenol and hydrofob amins have been studied by spectrophotometry. Extraction of mixed-ligand complexes is maximal at pH 3.8-5.3. The optimal conditions for the formation and extraction of mixed-ligand compounds have been found and the ratios of components in the complexes have been determined. A procedure has been developed for extraction –spectrophotometric determination Vanadium in soils and in oil and oil-products .

REFERENCES

1. Vijaya K., Kumar Reddy, Yamini P., Kishore Kumar R. and Venkateswarlu P.. International Journal of Chemical Engineering and Applied Sciences (2012), 2 (1): pp1-5
2. B. Krishna Priya, P. Subrahmanyam, J. Dilip Kumar and P. Chiranjeevi. E-Journal of Chemistry. (2006). Vol. 3, No.4, pp 286-297,
3. Richard, F. J. Dams, Jan Goossens and Luc Moens. (1995) Microchimica Acta 119 (3-4), pp 277-286.
4. Kawakubo, S., Tsuchiya, Y. and Iwatsuki, M. (1995) Anal. Chim. Acta 310, pp 501-507.
5. Mohamad, A. S. and Fawy, K. H. (2000). Mikrochim. Acta 134, pp 229-234.
6. Ensafi, A. A., Amini, M. K. and Mazloum, M. (1999) Anal. Lett. 32, pp 1927-1937.
7. Spinola Costa A.C., Leonardo S. G. Teixeira, Helena Valli Yaeger, and Sergio L. C. Ferreira. Spectrophotometric Determination of Vanadium (IV) in the Presence of Vanadium (V) Using Br-PADAP. (1998). Mikrochim Acta, 130, pp 41-45
8. Qiufen Hu, Yang G and Yin J, *Bull. Korean Chem. Soc.*, (2004), 25, p 263.
9. Kiran Kumar T. N and Revanasiddappa H D, *Iranian J. Chem. Soc.*, (2005), 2, p161.
10. Bosque-Sendra J. M, Valencia L C and Boudra S, *Fresenius' J. Anal. Chem.*, (1998), 360, 31.
11. Chauhan R S and Kakkar L R, *Bull. Chem. Soc. Japan*, (1992), 65, p 1033.
12. Iranpoor N, Maleki N, Razi S and Safavi A, *Talanta*, (1992), 39, p 281.
13. Miura J H, *Anal. Chim.*, 1990, 62, p 1424.
14. Murthy G V R, Reddy T S and Rao S B, *Analyst*, (1989), 114, p.493.
15. Taylor M J C, Marshall G D, Williams S J S, Vanstaden J F and Sailing C, *Anal. Chim. Acta*, (1996), 329, p 275.
16. Serrat F B and Morell G B, *Fresenius' J. Anal. Chim.*, (1994), 349, p 717.
17. Zucchi C, Forneris M, Martirez L, Oisina R and Marchevsky E, *Fresenius' J. Anal. Chem.*, (1998), 360, p 128.
18. Shigenori N, Tanaka E and Mizutani Y, *Talanta*, (2003), 61, p 203.



19. Verdizade, N. A. and Kuliev, K. A., Zh. Anal. Khim., (1998), vol. 53, no. 1, p 25 (J. Anal. Chem. (Engl. Transl.), vol.53, no. 1, p 19).
20. Verdizade, N. A., Amrakhov, T. I., Kuliev, K.A., and Zalov, A. Z., Zh. Anal. Khim., (1997), vol. 52, no.10,p.1042. J. Anal. Chem. (Engl.Transl.), vol. 52, no.10, p 944.
21. Verdizade, N. A., Kuliev, K. A., Zalov, A. Z., Efendieva, N. N., Alkhverdiev, M. A., and Iskenderov, M. G., Abstracts of papers, Vseros. Konf. "Khimicheskii analiz" (All-Russian Conf. on Chemical Analysis), Moscow, 2008, p 96.
22. Charlot, G., Les methods de la chimie analytique: anlyse quantitative minerale, Masson, 1961, 4th ed.
23. Marczenko, Z., Baltsejak, M. K. Metodi Spectrophotometrii v UF I vidimoy oblasti (Spectrophotometrically in the UV and visible regions in inorganic analysis). M. Binom. Laboratoriya znaniy. 2007. P 711.
24. Nazarenko, V. A., and Biryuk, E. A. (1967). A study of the chemistry of reactions of multivalent element ions with organic reagents. Zh. Anal. Khim., 22 (1), pp 57-64.
25. Nazarenko, V. A., Tr. Komis. Po analit. Khimii Akad. Nauk SSSR (Proc. Comissionon Analytical Chemistry of the USSR's Academy of Sciences), Moscow: Nauka, 1969, vol. 17, p 22.
26. Al'tshuler, S. A. and Kozyrev, B. M., Elektronnyi paramagnitnyi rezonans soedinenii promezhutochnykh grupp (Electron Paramagnetic Resonance of the Compounds of Intermediate Groups), Moscow: Nauka, 1972.
27. Bulatov, M. I. and Kalinkin, I. P., Prakticheskoe rukovodstvo po fotokolorimetriceskim I spektro-fotometriceskim metodam analiza (Practical Guide on Photocolorimetric and Spectropho-tometric Methods of Analysis), Moscow: Khimiya, 1972.
28. Nakamoto, K. IK-spectr KR neorganiceskikh I koordinatsionnikh soedinenii (Infrared and Raman Spectra of Inorganic and Coordination Compounds). Moscow: Mir, 1991. P 536.
29. Bellami, L. Infrakrasnie spectri slojnikh molecule (The infra-red spectra of complex molecules). Moscow: Mir, 1991. P 592.
30. Baranova, L. I., Orlova, E. S., and Tserkovnitskaya, I. A., Zavod. Lab., (1978), vol. 44, no. 2. p 142.