



ON A NEW SUBCLASS OF MULTIVALENT FUNCTIONS DEFINED BY HADAMARD PRODUCT

Waggas Galib Atshan*, Ali Hussein Battor** and Azhar Jawad Kadhim**

*Department of Mathematics, Collage of Computer Science and Mathematics
University of Al-Qadisiya, Diwaniya-Iraq

**Department of Mathematics, College of Education for Girls University of Kufa, Najaf-Iraq

**battor-ali@yahoo.com, *waggashnd@gmail.com, waggas-hnd@yahoo.com

**azharjawad91@yahoo.com

Abstract

In this paper, we introduce and study the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$ of multivalent functions in the open unit disk $U = \{z \in \mathbb{C} : |z| < 1\}$, which are defined by the convolution (or Hadamard product). We give some properties, coefficient inequality, closure theorems, neighborhoods of the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$, partial sums, weighted mean theorem, convolution, distortion and growth bounds.

2015 **Mathematics Subject Classification:** 30C45; 30C50

Keywords: Multivalent Function; Closure theorem; neighborhood; Partial Sums; Convolution; Distortion bounds.



Council for Innovative Research

Peer Review Research Publishing System

Journal: JOURNAL OF ADVANCES IN MATHEMATICS

Vol .10, No.5

www.cirjam.com , editorjam@gmail.com



INTRODUCTION

Let M_p be denote the class of all functions of the form:

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n, \quad (p \in N = \{1, 2, \dots\}), \quad (1.1)$$

which are analytic and multivalent in the open unit disk $U = \{z \in \mathbb{C}: |z| < 1\}$.

Let M_p^* be denote the subclass of M_p consisting of functions of the form:

$$f(z) = z^p - \sum_{n=p+1}^{\infty} a_n z^n, \quad (a_n \geq 0, p \in N). \quad (1.2)$$

For the function $f \in M_p^*$ given by (1.2) and $g \in M_p^*$ defined by

$$g(z) = z^p - \sum_{n=p+1}^{\infty} b_n z^n, \quad (b_n \geq 0, p \in N). \quad (1.3)$$

We define the convolution (or Hadamard product) of f and g by

$$(f * g)(z) = z^p - \sum_{n=p+1}^{\infty} a_n b_n z^n = (g * f)(z). \quad (1.4)$$

Definition(1): For $0 \leq \lambda < \frac{1}{2}, -1 \leq \alpha < 0, 0 \leq \mu < 1$ and $-\frac{1}{3} < \zeta \leq 0, p \in N$, a function $f \in M_p^*$ is said to be in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$ if it satisfies the condition:

$$Re \left\{ \frac{z^2((f * g)(z))'' - pz((f * g)(z))'}{\lambda z^2((f * g)(z))'' - (\alpha p + [(1 - \mu)(1 - \alpha)\zeta])z((f * g)(z))'} \right\} > \beta. \quad (1.5)$$

Some authors studied multivalent functions for another classes, like, ([2], [3], [4],[5]).

2.Coefficient bounds:

Lemma(1)[1]: Let $w = (u + iv)$ is a complex number, then $Re(w) > \beta$ if and only if $|w - (p - \beta)| < |w + (p + \beta)|$, where $\beta \geq 0$.

Theorem(1): Let $f \in M_p^*$. Then $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$ if and only if

$$\sum_{n=p+1}^{\infty} n[\lambda(n-1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]] a_n b_n \leq p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]], \quad (2.1)$$

where

$$0 \leq \lambda < \frac{1}{2}, -1 \leq \alpha < 0, 0 \leq \mu < 1, -\frac{1}{3} < \zeta \leq 0 \text{ and } p \in N.$$

The result is sharp for the function

$$f(z) = z^p - \frac{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]}{n[\lambda(n-1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]} z^n.$$

Proof: Suppose that the inequalities (2.1) holds and let $|z| = 1$, in view of

(1.5), we need to prove that $Re(w) > \beta$, where

$$w = \frac{z^2((f * g)(z))'' - pz((f * g)(z))'}{\lambda z^2((f * g)(z))'' - (\alpha p + [(1 - \mu)(1 - \alpha)\zeta])z((f * g)(z))'}$$



$$= \frac{-p - \sum_{n=p+1}^{\infty} (n^2 + n(p-1))a_n b_n z^{n-p}}{p(p(\lambda - \alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]) - \sum_{n=p+1}^{\infty} n[\lambda(n-1) - \alpha p - ((1-\mu)(1-\alpha)\zeta)]a_n b_n z^{n-p}}$$

$$= \frac{A(z)}{B(z)}$$

By Lemma (1), it suffice to show that

$$|A(z) - (p + \beta)B(z)| - |A(z) + (p - \beta)B(z)| \leq 0, (0 \leq \beta < p).$$

Therefore, we obtain

$$\begin{aligned} & |A(z) - (p + \beta)B(z)| - |A(z) + (p - \beta)B(z)| \\ & \leq -(p + \beta)p[p(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)] \\ & \quad + (p + \beta) \sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]a_n b_n z^{n-p} \\ & \quad - (p - \beta)p[p(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)] \\ & \quad + (p - \beta) \sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]a_n b_n z^{n-p} \\ & = -2p^2[p(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)] + 2p \sum_{n=p+1}^{\infty} [\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]a_n b_n \leq 0, \end{aligned}$$

by hypothesis. Then by maximum modulus theorem, we have $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$.

Conversely, assume

$$\begin{aligned} & \operatorname{Re} \left\{ \frac{z^2((f * g)(z))'' - pz((f * g)(z))'}{\lambda z^2((f * g)(z))'' - (\alpha p + [(1 - \mu)(1 - \alpha)\zeta])z((f * g)(z))'} \right\} \\ & = \operatorname{Re} \left\{ \frac{-p - \sum_{n=p+1}^{\infty} (n^2 + n(p-1))a_n b_n z^{n-p}}{p(p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]) - \sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]a_n b_n z^{n-p}} \right\} \\ & > 1. \end{aligned} \quad (2.2)$$

We choose the value of z on the real axis let $z \rightarrow 1^-$ through real values, we can write (2.2) as

$$\sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]a_n b_n \leq p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]].$$

Finally, sharpness follows if we take

$$f(z) = z^p - \frac{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]}{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]} z^n, n \geq p + 1. \quad (2.3)$$

Corollary(1): Let $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$. Then

$$a_n \leq \frac{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]}{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]} b_n, n \geq p + 1. \quad (2.4)$$

3. Extreme Points:

In the following theorem, we obtain extreme points for the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$.

Theorem(2): Let $f_p(z) = z^p$ and

$$f_n(z) = z^p - \frac{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]}{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]} z^n, n \geq p + 1.$$

Then $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$ if and only if it can be expressed in the form



$$f(z) = \sum_{n=p}^{\infty} \theta_n f_n(z),$$

where $\theta_n \geq 0$ and

$$\sum_{n=p}^{\infty} \theta_n = 1.$$

Proof: Assume that

$$f(z) = \sum_{n=p}^{\infty} \theta_n f_n(z),$$

hence we get

$$f(z) = z^p - \sum_{n=p+1}^{\infty} \theta_n \frac{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]}{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]} z^n.$$

Now, $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$, since

$$\begin{aligned} & \sum_{n=p+1}^{\infty} \frac{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]b_n}{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]} \times \frac{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]\theta_n}{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]b_n} \\ &= \sum_{n=p+1}^{\infty} \theta_n = 1 - \theta_1 \leq 1. \end{aligned}$$

Conversely, suppose that $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$. Then we show that f can be written in the form $\sum_{n=p}^{\infty} \theta_n f_n(z)$.

Now $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$ implies from Theorem (1)

$$a_n \leq \frac{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]}{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]} b_n, n \geq p + 1.$$

Setting

$$\theta_n = \frac{n[\lambda(n - 1) - \alpha p - ((1 - \mu)(1 - \alpha)\zeta)]b_n}{p[(\lambda - \alpha) - \lambda - ((1 - \mu)(1 - \alpha)\zeta)]} a_n$$

and

$$\theta_p = 1 - \sum_{n=p+1}^{\infty} \theta_n.$$

We obtain

$$f(z) = \sum_{n=p}^{\infty} \theta_n f_n(z).$$

4. Closure Theorem:

Now, we shall prove the closure theorem of the functions in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$.

Theorem(3): Let $f_r \in M_p^*(\lambda, \alpha, \mu, \zeta, p), r = 1, 2, \dots, \ell$. Then

$$h(z) = \sum_{r=1}^{\ell} c_r f_r(z) \in M_p^*(\lambda, \alpha, \mu, \zeta, p).$$

For $f_r(z) = \sum_{n=p+1}^{\infty} a_{n,r} z^n$, where $\sum_{r=1}^{\ell} c_r = 1$.



Proof:

$$h(z) = \sum_{r=1}^{\ell} c_r f_r(z)$$

$$= z^p - \sum_{n=p+1}^{\infty} \sum_{r=1}^{\ell} c_r a_{n,r} z^n = z^p - \sum_{n=p+1}^{\infty} e_n z^n,$$

where $e_n = \sum_{r=1}^{\ell} c_r a_{n,r}$. Thus $h(z) \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$ if

$$\sum_{n=p+1}^{\infty} \frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]] b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]} e_n \leq 1,$$

that is, if

$$\sum_{n=p+1}^{\infty} \sum_{r=1}^{\ell} \frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]] b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]} c_r a_{n,r}$$

$$\sum_{r=1}^{\ell} c_r \sum_{n=p+1}^{\infty} \frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]] b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]} a_{n,r} \leq \sum_{r=1}^{\ell} c_r = 1.$$

5. Convolution:

In the following theorem, we obtain the convolution result of functions belong to the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$.

Theorem (4): Let the functions $f_j(z)$, ($j = 1, 2$) defined by

$$f_j(z) = z^p - \sum_{n=p+1}^{\infty} a_{n,j} z^n, (j = 1, 2)$$

be in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$. Then the function

$$T(z) = z^p - \sum_{n=p+1}^{\infty} (a_{n,1}^2 + a_{n,2}^2) z^n,$$

also belong to the class $M_p^*(\lambda, \alpha, \mu, \epsilon, p)$, where

$$\epsilon \geq \frac{A}{B},$$

where

$$A = p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]^2 [\lambda(n-1) - \alpha p] + n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]]^2 b_n [\lambda - p(\lambda-\alpha)],$$

$$\text{and } B = [(1-\mu)(1-\alpha)] [p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]^2 - n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]]^2 b_n].$$

Proof: From Theorem (1), we have

$$\sum_{n=p+1}^{\infty} \left(\frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]] b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]} \right)^2 a_{n,j}^2 \leq \left(\sum_{n=p+1}^{\infty} \frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]] b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]} a_{n,j} \right)^2 \leq 1,$$

it follows that

$$\sum_{n=p+1}^{\infty} \frac{1}{2} \left(\frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]] b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]} \right)^2 (a_{n,1}^2 + a_{n,2}^2) \leq 1.$$

But $T \in M_p^*(\lambda, \alpha, \mu, \epsilon, p)$ if and only if



$$\sum_{n=p+1}^{\infty} \frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\epsilon]]b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\epsilon]]} (a_{n,1}^2 + a_{n,2}^2) \leq 1, \quad (5.1)$$

the inequality (5.1) will be satisfied if

$$\frac{n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\epsilon]]b_n}{p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\epsilon]]} \leq \frac{n^2[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]]^2 b_n^2}{p^2[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]^2}, \quad (n \geq p+1)$$

so that

$$\epsilon \geq \frac{A}{B},$$

where

$$A = p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]^2 [\lambda(n-1) - \alpha p] + n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]]^2 b_n [\lambda - p(\lambda-\alpha)],$$

$$\text{and } B = [(1-\mu)(1-\alpha)] [p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]^2 - n[\lambda(n-1) - \alpha p - [(1-\mu)(1-\alpha)\zeta]]^2 b_n].$$

This completes the proof.

6. Neighborhoods:

Following the earlier works on neighborhoods of analytic functions by Goodman [6] and Ruscheweyh [7], we begin by introducing here the δ -neighborhood of a function $f \in M_p^*$ of the form (1.2) by means of the definition below:-

$$N_\delta(f) = \left\{ g \in M_p^*; g(z) = z^p - \sum_{n=p+1}^{\infty} b_n z^n \text{ and } \sum_{n=p+1}^{\infty} n|a_n - b_n| \leq \delta, 0 \leq \delta < 1 \right\}. \quad (6.1)$$

Particularly for the identity function $\vartheta(z) = z^p$, we have

$$N_\delta(z) = \left\{ g \in M_p^*; g(z) = z^p - \sum_{n=p+1}^{\infty} b_n z^n \text{ and } \sum_{n=p+1}^{\infty} n|b_n| \leq \delta \right\}.$$

Definition(2): A function $f \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$ is said to be in the class $M_{p,\vartheta}^*(\lambda, \alpha, \mu, \zeta, p)$ if there exists function $g \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$ such that

$$\left| \frac{f(z)}{g(z)} - 1 \right| < 1 - \vartheta, \quad (z \in U, 0 \leq \vartheta < 1).$$

Theorem(5): If $g \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$ and

$$\vartheta = 1 - \frac{\delta[(p+1)(p(\lambda-\alpha) - [(1-\mu)(1-\alpha)\zeta])]a_{p+1}}{(p+1)(p(\lambda-\alpha) - [(1-\mu)(1-\alpha)\zeta])a_{p+1} - p[p(\lambda-\alpha) - \lambda - [(1-\mu)(1-\alpha)\zeta]]}. \quad (6.2)$$

Then $N_\delta(g) \subset M_{p,\vartheta}^*(\lambda, \alpha, \mu, \zeta, p)$.

Proof: Let $f \in N_\delta(g)$. Then we find from (6.2) that

$$\sum_{n=p+1}^{\infty} n|a_n - b_n| \leq \delta,$$

which implies the coefficient inequality

$$\sum_{n=p+1}^{\infty} |a_n - b_n| \leq \delta, \quad (n \geq p+1). \quad (6.3)$$



Since $g \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$, then by using Theorem (1)

$$\sum_{n=p+1}^{\infty} b_n \leq \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]a_{p+1}} \tag{6.4}$$

So that

$$\left| \frac{f(z)}{g(z)} - 1 \right| \leq \frac{\sum_{n=p+1}^{\infty} |a_n - b_n|}{1 - \sum_{n=p+1}^{\infty} b_n} \leq \frac{\delta[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]a_{p+1}}{(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])a_{p+1} - p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}$$

$$= 1 - \vartheta.$$

Hence by definition (2) $f \in M_{p,\vartheta}^*(\lambda, \alpha, \mu, \zeta, p)$ for ϑ given by (6.2). This complete the proof.

Theorem(6): Let $f(z) \in M_p^*$ be given by (1.2) and define the partial sums

$s_1(z)$ and $s_v(z)$ by

$$s_1(z) = z^p$$

$$s_v(z) = z^p + \sum_{n=p+1}^{p+v-1} a_n z^n, \quad v > p + 1 \tag{6.5}$$

suppose also that

$$\sum_{n=p+1}^{\infty} d_n a_n \leq 1,$$

$$d_n = \left(\frac{n[\lambda(n - 1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]]b_n}{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]} \right). \tag{6.6}$$

Thus, we have

$$Re \left\{ \frac{f(z)}{s_v(z)} \right\} > 1 - \frac{1}{d_n} \tag{6.7}$$

and

$$Re \left\{ \frac{s_v(z)}{f(z)} \right\} > 1 - \frac{d_n}{1 + d_n}. \tag{6.8}$$

Each of the bounds in (6.7) and (6.8) is the best possibility for $p \in N$.

Proof: For the coefficients d_n given by (6.6), it is difficult to verify that

$$d_{n+1} > d_n > 1, \quad n \geq p + 1.$$

Therefore, by using the hypothesis (6.5), we have

$$\sum_{n=p+1}^{p+v-1} a_n + d_n \sum_{n=p+v}^{\infty} a_n \leq \sum_{n=p+1}^{\infty} d_n a_n \leq 1. \tag{6.9}$$

By setting

$$g_1(z) = d_n \left(\frac{f(z)}{s_v(z)} - \left(1 - \frac{1}{d_n} \right) \right) = 1 + \frac{d_n \sum_{n=p+v}^{\infty} a_n z^{n-p}}{1 + \sum_{n=p+1}^{p+v-1} a_n z^{n-p}} \tag{6.10}$$

and applying (6.9), we find that

$$\left| \frac{g_1(z) - 1}{g_1(z) + 1} \right| \leq \frac{d_n \sum_{n=p+v}^{\infty} a_n}{2 - 2 \sum_{n=p+1}^{p+v-1} a_n - d_n \sum_{n=p+v}^{\infty} a_n} \leq 1.$$



This proves (6.7). Therefore, $\operatorname{Re}(g_1(z)) > 0$ and we obtain that

$$\operatorname{Re}\left\{\frac{f(z)}{s_v(z)}\right\} > 1 - \frac{1}{d_n}.$$

Now, in the same manner, we can prove the assertion (6.8), by setting

$$g_2(z) = (1 + d_n)\left(\frac{s_v(z)}{f(z)} - \frac{d_n}{1 + d_n}\right).$$

This complete the proof.

7. Weighted mean:

Definition(3): Let f and g be in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$. Then, the weighted mean E_q of f and g is given by

$$E_q(z) = \frac{1}{2}[(1 - q)f(z) + (1 + q)g(z)], \quad 0 < q < 1.$$

Theorem(7): Let f and g be in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$. Then, the weighted mean of f and g is also in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$.

Proof: By definition (3), we have

$$\begin{aligned} E_q(z) &= \frac{1}{2}[(1 - q)f(z) + (1 + q)g(z)] \\ &= \frac{1}{2}\left[(1 - q)\left(z^p - \sum_{n=p+1}^{\infty} a_n z^n\right) + (1 + q)\left(z^p - \sum_{n=p+1}^{\infty} b_n z^n\right)\right] \\ &= z^p - \sum_{n=p+1}^{\infty} \frac{1}{2}((1 - q)a_n + (1 + q)b_n)z^n. \end{aligned}$$

Since f and g are in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$ so by Theorem (1), we get

$$\sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]] a_n \leq p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]$$

and

$$\sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]] b_n \leq p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]].$$

Hence,

$$\begin{aligned} &\sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]] \left(\frac{1}{2}(1 - q)a_n + \frac{1}{2}(1 + q)b_n\right) \\ &\frac{1}{2}(1 - q) \sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]] a_n + \frac{1}{2}(1 + q) \sum_{n=p+1}^{\infty} n[\lambda(n - 1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]] b_n \\ &\leq \frac{1}{2}(1 - q)p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]] + \frac{1}{2}(1 + q)p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]] \\ &= p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]. \end{aligned}$$

This shows $E_q \in M_p^*(\lambda, \alpha, \mu, \zeta, p)$.

8. Distortion and growth bounds:

In the following theorems, we prove distortion and growth bounds.



Theorem(8): Let the function f defined by (1.2) be in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$. Then

$$\begin{aligned} r^p - \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]b_{p+1}} r^{p+1} &\leq |f(z)| \\ &\leq r^p + \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]b_{p+1}} r^{p+1}, \\ 0 < |z| = r < 1. \end{aligned} \quad (8.1)$$

the equality in (8.1) is attained by the function f given by

$$f(z) = z^p - \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]b_{p+1}} z^{p+1},$$

Proof: Since the function f defined by (1.2) in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$ we have from Theorem (1),

$$\sum_{n=p+1}^{\infty} a_n \leq \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]b_{p+1}}.$$

Thus

$$|f(z)| \leq |z|^p + \sum_{n=p+1}^{\infty} a_n |z|^n = r^p + r^{p+1} \sum_{n=p+1}^{\infty} a_n \leq r^p + \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]b_{p+1}} r^{p+1}.$$

Similarly

$$|f(z)| \geq |z|^p - \sum_{n=p+1}^{\infty} a_n |z|^n = r^p - r^{p+1} \sum_{n=p+1}^{\infty} a_n \geq r^p - \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]b_{p+1}} r^{p+1}.$$

Theorem(9): Let the function f defined by (1.2) in the class $M_p^*(\lambda, \alpha, \mu, \zeta, p)$,

$$(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])b_{p+1} \leq n[\lambda(n - 1) - \alpha p - [(1 - \mu)(1 - \alpha)\zeta]]b_n$$

Then

$$pr^{p-1} - \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta]]b_{p+1}} r^p \leq |f'(z)| \leq pr^{p-1} + \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta]]b_{p+1}} r^p, 0 < |z| = r < 1, \quad (8.2)$$

the equality in (8.2) is attained by the function f given by

$$f(z) = z^p - \frac{p[p(\lambda - \alpha) - \lambda - [(1 - \mu)(1 - \alpha)\zeta]]}{[(p + 1)(p(\lambda - \alpha) - [(1 - \mu)(1 - \alpha)\zeta])]b_{p+1}} z^{p+1},$$

Proof: Theorem (9) can be proved easily by the similar steps of Theorem (8).

References

- [1] E. S. Aqlan , **Some Problems Connected with Geometric Function Theory** , Ph. D. thesis (2004), Pune University, Pune .
- [2]W. G. Atshan, **On a certain class of multivalent functions defined by Hadamard product**, Journal of Asian Scientific Research, 3(9)(2013), 891-902.
- [3] W. G. Atshan, A. H. Battor and A. A. Abdulridha, **Some Aspects of new subclass of multivalent functions defined by Integral operator**, 2nd International Conference of Mathematics and its Application 23-24 October (2013).
- [4] W. G. Atshan, A. H. Battor and A. M. Dereush, **On a new class of multivalent functions with negative coefficients defined by Hadamard product involving a linear operator**, American Journal of Mathematics and statistics, (USA), 4(3)(2014), 147-155.
- [5] W. G. Atshan, H. J. Mustafa and E. K. Mouajeeb, **Subclass of multivalent functions defined by Hadamard product involving a linear operator**, International Journal of Math. Analysis, 7(24)(2013), 1193-1206.



- [6] A. W. Goodman, **Univalent functions and analytic curves**, Proc. Amer. Math. Soc., 8(1975), 598-601.
[7] S. Ruscheweyh, **Neighborhoods of univalent functions**, Proc. Amer. Math. Soc., 81 (1981), 521-527.

