



## PARTITION OF MEASURABLE SETS

Otanga L. Olwamba, Owino M. Oduor, Aywa S. Omukunda

<sup>1</sup> Department of Mathematics,  
Masinde Muliro University of Science and Technology P.O. Box 190-50100, Kakamega, Kenya  
leviotanga@gmail.com

<sup>2</sup> Department of Mathematics and Computer Science, University of Kabianga P.O Box 2030-20200, Kericho,  
Kenya  
morricearaka@yahoo.com

<sup>3</sup> Department of Mathematics, Kibabii University College P.O Box 1699-50200, Bungoma, Kenya  
shemaywa@gmail.com

### Abstract

The theory of vector measure has attracted much interest among researchers in the recent past. Available results show that measurability concepts of the Lebesgue measure have been used to partition subsets of the real line into disjoint sets of finite measure. In this paper we partition measurable sets in  $\mathfrak{R}^n$  for  $n \geq 3$  into disjoint sets of finite dimension.

**AMS Subject Classification:** Primary 20E45, 20H30; Secondary 28B15, 28B20

**Key Words and Phrases:** Partition; Measurable cover; Extension procedures; countable additivity.



## Council for Innovative Research

Peer Review Research Publishing System

Journal: JOURNAL OF ADVANCES IN MATHEMATICS

Vol .10, No.8

[www.cirjam.com](http://www.cirjam.com) , [editorjam@gmail.com](mailto:editorjam@gmail.com)



## 1 INTRODUCTION

In this article, measurable cover estimate technique is applied to partition measurable sets. The utility of properties such as extension, countable additivity and contraction of the projective tensor product of vector measure duality is demonstrated. The process of partition requires that we vary all countable coverings of sets in a ring by sets in a  $\sigma$ -ring. Sequences of monotonically increasing (or decreasing) sets and integrability concepts of strictly increasing (or decreasing) functions are considered. For notations and basic concepts used in this paper, reference may be made to [1, 2, 3, 4, 5, 6]

## 2 Basic Concepts

### 2.1 Projective tensor product vector measure duality

Let  $X_1, \dots, X_n$  and  $Z$  be real Banach spaces with  $\Phi: \prod_{i=1}^n X_i \rightarrow Z$  being a continuous linear function. If  $\mu_1: R_1 \rightarrow X_1, \dots, \mu_n: R_n \rightarrow X_n$  are countably additive vector measures, then the product  $\prod_{i=1}^n \mu_i$  is a countably additive vector measure defined on the ring  $\prod_{i=1}^n R_i$  generated by sets of the form  $E_1 \times \dots \times E_n$ . Let  $(G(R_1), \dots, G(R_n))$  be a set of  $\sigma$ -rings generated by rings  $R_1, \dots, R_n$  respectively. If the extension of the vector measure  $\prod_{i=1}^n \mu_i: \prod_{i=1}^n R_i \rightarrow \prod_{i=1}^n X_i$  to a vector measure  $\prod_{i=1}^n \mu_i^*: \prod_{i=1}^n G(R_i) \rightarrow \prod_{i=1}^n X_i$  coincide with respect to a linear function  $\Phi: \prod_{i=1}^n X_i \rightarrow Z$ , where  $Z = \Phi(\mu_1^*(E_1), \dots, \mu_n^*(E_n))$  for  $\mu_i^*(E_i) \in X_i, 1 \leq i \leq n$  and  $\prod_{i=1}^n X_i$  is a Banach space, then  $(\prod_{i=1}^n)_{\Phi} < \mu_i^*(E_i), Z' >$  is called the projective tensor product of vector measure duality between  $Z$  and its dual space  $Z'$ .

### 2.2 3 D section of a measurable set

Fix  $p_{n-2}$  for  $n \in \mathbb{N}$  and  $n \geq 3$ . The set  $(\prod_{i=1}^n E_i)^{p_{n-2}} = E_{n-2} \times E_{n-1} \times E_n$  is called a three dimensional section of a measurable set  $\prod_{i=1}^n E_i = N(f)$  such that  $\iiint f \delta(\prod_{i=1}^n)_{\Phi} < \mu_i^*(E_i), Z' > < \infty$ .

## 3 Results

**Proposition 1** Let  $f: \prod_{i=1}^n G(R_i) \rightarrow \prod_{i=1}^n X_i$  be an integrable function with respect to  $(\prod_{i=1}^n)_{\Phi} < \mu_i^*, Z' >$ . Suppose the set  $((e_{n-2}, e_{n-1}, e_n): f(e_{n-2}, e_{n-1}, e_n) > P)$  for  $P > 0$ , is measurably covered by  $N(f)$ . If  $(\prod_{i=1}^n E_i)_{e_{n-2}}$  is a countable partition of a set  $((e_{n-2}, e_{n-1}, e_n): f(e_{n-2}, e_{n-1}, e_n) > P)_{e_{n-2}}$  for a fixed  $e_{n-2} \in E_{n-2}$  such that  $\iiint f \delta(\prod_{i=1}^n)_{\Phi} < \mu_i^*(E_i), Z' > < M$  for  $M > 0$ , then  $< T_{(\prod_{i=1}^n)_{\Phi} \mu_i^*(E_i)}(f), Z' > = 0$

**Proof.** Let  $N(f) = ((e_{n-2}, e_{n-1}, e_n): f(e_{n-2}, e_{n-1}, e_n) \neq 0)$ . It can be deduced from [2, p.18] that  $(\prod_{i=1}^n E_i)_{e_{n-2}} = \prod_{i=1}^n E_i$  for a fixed  $e_{n-2} \in E_{n-2}$ . Since by hypothesis  $((e_{n-2}, e_{n-1}, e_n): f(e_{n-2}, e_{n-1}, e_n) > P) \ominus N(f)$  and  $(\prod_{i=1}^n E_i)_{e_{n-2}}$  is a countable partition of  $((e_{n-2}, e_{n-1}, e_n): f(e_{n-2}, e_{n-1}, e_n) > P)_{e_{n-2}}$ , it follows that  $(\prod_{i=1}^n E_i)_{e_{n-2}} = \prod_{i=1}^n E_i \ominus (N(f))_{e_{n-2}} \Rightarrow ((e_{n-2}, e_{n-1}, e_n): f(e_{n-2}, e_{n-1}, e_n) > P)_{e_{n-2}} = \bigcup_{k=1}^{\infty} \prod_{i=1}^n E_{k_i}$  for a countable partition  $\prod_{i=1}^n E_{k_i}$  of  $\prod_{i=1}^n E_i$ . In this case,  $\prod_{i=1}^n E_i$  can be written uniquely as a countable union of disjoint sets (see [3 p.5]). Now,  $\bigcup_{k=1}^{\infty} \prod_{i=1}^n E_{k_i} = \prod_{i=1}^n E_i \Rightarrow P \chi_{\prod_{i=1}^n E_i} \leq \chi_{\prod_{i=1}^n E_i} f$ . Let  $(N(f))_{e_{n-2}} = \prod_{i=1}^n C_i$  be a set in a  $\sigma$ -ring  $\prod_{i=1}^n G(R_i)$ . It follows that  $\prod_{i=1}^n E_i \subset \prod_{i=1}^n C_i \Rightarrow \prod_{i=1}^n E_i \ominus \prod_{i=1}^n C_i$ . Let  $\mu_{i_{C_i}}^*$  be a vector measure defined on  $G(R_i)$  for  $i = n-1, n$ .

. By extension procedures and integral representation of the contraction of  $\mu_{n-1} \times \mu_n$  as illustrated in [6], we have  $P < T_{(\prod_{i=1}^n)_{\Phi} \mu_i^*(C_i \cap E_i)}(f), Z' > \leq \iiint f \delta(\prod_{i=1}^n)_{\Phi} < \mu_i^*(E_i), Z' > < M P < T_{(\prod_{i=1}^n)_{\Phi} \mu_i^*(C_i \cap E_i)}(f), Z' > \leq \iiint f \delta(\prod_{i=1}^n)_{\Phi} < \mu_i^*(C_i \cap E_i), Z' > < M \Rightarrow < T_{(\prod_{i=1}^n)_{\Phi} \mu_i^*(C_i \cap E_i)}(f), Z' > < M/P$  But  $E_i \subset C_i$  for



$1 = n - 1, n$ . Therefore,  $\langle T_{(\Pi_{i=n-1}^n)_\Phi \mu_i^*(E_i)}(f), Z' \rangle = M/P$ . As  $P \rightarrow \infty$ ,  $\langle T_{(\Pi_{i=n-1}^n)_\Phi \mu_i^*(E_i)}(f), Z' \rangle = 0$

**Proposition 2** Let  $f$  and  $f_n (n = 1, 2, \dots)$  for  $f_n \uparrow$  be integral functions with respect to  $(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*, Z' \rangle$  such that  $N(f_n)$  is measurably covered by  $N(f)$ . If  $E_i = N(g_n) \subset N(f) - N(f_n)$  for  $(n = 1, 2, \dots)$  and  $i = n - 2, n - 1, n$ , we have  $\iiint g_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i), Z' \rangle < \varepsilon$ , for  $\varepsilon > 0$

**Proof.** Let  $N(f_n) = ((e_{n-2}, e_{n-1}, e_n) : f_n(e_{n-1}, e_{n-1}, e_n) \neq 0)$  and  $N(f) = ((e_{n-2}, e_{n-1}, e_n) : f(e_{n-2}, e_{n-1}, e_n) \neq 0)$  be measurable sets of finite measure. Since  $N(g_n) \subset N(f) - N(f_n)$  for each  $n$ , it follows that  $g_n \downarrow 0$  for each  $n$ . Let  $E_{n-2} = N(g_{n-2})$  and  $A_{k_i}$  be a countable partition of  $E_i$  for  $i = n - 2, n - 1, n$ . Now,  $A_{k_i} \uparrow E_i$  for  $i = n - 2, n - 1, n$  (see [3 p.20]). Suppose  $N(g_{n-1}) = E_i - \cup_{k=1}^\infty A_{k_i}$  for  $i = n - 2, n - 1, n$ . It follows that  $N(g_{n-1}) \downarrow \emptyset$ . Since  $g_{n-1}$  is integrable with respect to  $\Pi_{i=n-1}^n \langle \mu_i^*, Z' \rangle$  (see [5]),  $\Rightarrow \sum_{k=1}^\infty \iiint g_{n-1} \delta(\Pi_{i=n-1}^n)_\Phi \langle \mu_i^*(E_i - A_{k_i}), Z' \rangle = 0$ . Let us vary the partition  $A_{k_{o_i}}^j$  of  $A_{k_i}$  such that  $\Pi_{i=n-2}^n A_{k_i} = \cup_{j=1}^\infty \cup_{k_o=1}^\infty A_{k_{o_i}}^j$ . It follows that  $\sum_{j=1}^\infty \sum_{k_o=1}^\infty \iiint g_{n-1} \delta(\Pi_{i=n-1}^n)_\Phi \langle \mu_i^*(E_i - A_{k_{o_i}}^j), Z' \rangle < \varepsilon$ . Let  $A_i = \cup_{j=1}^\infty \cup_{k_o=1}^\infty A_{k_{o_i}}^j \Rightarrow \iiint g_{n-1} \delta(\Pi_{i=n-1}^n)_\Phi \langle \mu_i^*(E_i - A_i), Z' \rangle < \varepsilon$ . As shown in [3, p. 21], if we partition  $E_i$  into disjoint sets  $E_i - A_i$  and  $A_i$  for  $i = n - 2, n - 1, n$  and apply the additivity property of multiple integral, we have  $\iiint g_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i), Z' \rangle = \iiint g_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(A_i), Z' \rangle$ . Since  $A_i$  is a countable partition of  $E_i$ , then  $A_i \uparrow E_i$  and  $A_i \subset E_i$  for each  $i$ . Since  $E_i = N(g_n)$ ,  $g_n \downarrow 0$  for  $i = n - 2, n - 1, n$ , and  $n = 1, 2, \dots \Rightarrow \iiint g_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(A_i), Z' \rangle = 0$ , it follows that  $\iiint g_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i), Z' \rangle < \varepsilon$ .

**Proposition 3** Let  $\Pi_{i=n-2}^n E_i$  be measurably covered by  $N(f_n)$ . If  $\Pi_{i=n-2}^n C_i$  is a section of  $\Pi_{i=n-2}^n E_i$  such that  $C_{k_i} \downarrow \emptyset$  for every countable partition  $C_{k_i}$  of  $C_i$  for  $i = n - 2, n - 1, n$ , then

$$\iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i), Z' \rangle = \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i - C_i), Z' \rangle + \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_i - C_{k_i}), Z' \rangle$$

**Proof.** Choose  $\beta > 0$  such that  $\Pi_{i=n-2}^n C_i = ((x_{n-2}, x_{n-1}, x_n) : f(x_{n-2}, x_{n-1}, x_n) > \beta)$ . Since  $\Pi_{i=n-2}^n C_i$  is a section of  $\Pi_{i=n-2}^n E_i$ , then  $C_i \uparrow E_i$  for  $i = n - 2, n - 1, n$ . Let  $\iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i - C_i), Z' \rangle < \varepsilon$  for all  $n$ . Suppose we partition each  $E_i$  into disjoint sets  $E_i - C_i$  and  $C_i$  for  $i = n - 2, n - 1, n$ . Since  $\Pi_{i=n-2}^n E_i \ominus N(f_n)$

$$\text{, it follows that } \chi_{\Pi_{i=n-2}^n E_i} f_n = \chi_{\Pi_{i=n-2}^n (E_i - C_i)} f_n + \chi_{\Pi_{i=n-2}^n C_i} f_n \Rightarrow \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i), Z' \rangle = \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i - C_i), Z' \rangle + \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_i), Z' \rangle.$$

Since  $C_{k_i}$  is a countable partition of  $C_i$ , then  $C_i$  can be expressed as a union of disjoint measurable sets  $E_i - C_{k_i}$  and  $C_{k_i}$ . Therefore,

$$\iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_i), Z' \rangle = \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_i - C_{k_i}), Z' \rangle + \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_{k_i}), Z' \rangle.$$

By hypothesis,  $C_{k_i} \downarrow \emptyset$ . It follows that  $\iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_{k_i}), Z' \rangle = 0$ . (see [1, p. 2]).

$$\iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_i), Z' \rangle = \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_i - C_{k_i}), Z' \rangle. \text{ From the results above, we have } \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i), Z' \rangle = \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(E_i - C_i), Z' \rangle + \iiint f_n \delta(\Pi_{i=n-2}^n)_\Phi \langle \mu_i^*(C_i - C_{k_i}), Z' \rangle$$



**Proposition 4** Let  $f$  be an integrable function. Suppose  $\Pi_{i=n-2}^n E_{k_i}$  is measurably covered by  $\Pi_{i=n-2}^n(N(f))_i$  in  $\Pi_{i=n-2}^n G(R_i)$  such that  $\int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(N(f))_i, Z' > < M$  for  $M > 0$ . If  $\Pi_{i=n-2}^n E_{k_i}$  is a countable partition of  $\Pi_{i=n-2}^n E_i$  in  $\Pi_{i=n-2}^n R_i$  such that  $\Pi_{i=n-2}^n \mu_i(E_{k_i}) < M/\varepsilon$ , for  $\varepsilon > 0$ , then  $\Pi_{i=n-2}^n E_i \ominus \Pi_{i=n-2}^n(N(f))_i$

**Proof.** Let  $\Pi_{i=n-2}^n(N(f))_i = ((a, b, c) : f(a, b, c) \neq 0) \in \Pi_{i=n-2}^n G(R_i)$ . Suppose

$\Pi_{i=n-2}^n E_{k_i} = ((a, b, c) : f_n(a, b, c) > \varepsilon)$  and  $\Pi_{i=n-2}^n E_i = ((a, b, c) : f(a, b, c) > \varepsilon)$  are sets in  $\Pi_{i=n-2}^n R_i$ . Since  $\Pi_{i=n-2}^n E_{k_i}$  is a countable partition of  $\Pi_{i=n-2}^n E_i$ , we have  $E_{k_i} \uparrow E_i$  for  $i = n-2, n-1, n$  and each  $k \in \aleph$  (see [3 p.20]). So,  $\Pi_{i=n-2}^n E_i = \cup_{k=1}^\infty \Pi_{i=n-2}^n E_{k_i}$ . Hence,  $\varepsilon \chi_{\Pi_{i=n-2}^n E_{k_i}} \leq \chi_{\Pi_{i=n-2}^n E_{k_i}} f_n$ . Let  $\mu_{i=N(f)_i}^*$  be a vector measure

defined on  $G(R_i)$  for  $i = n-2, n-1, n$ . Since  $\Pi_{i=n-2}^n E_{k_i}$  is measurably covered by  $\Pi_{i=n-2}^n(N(f))_i$  for  $i = n-2, n-1, n$ , we have  $E_{k_i} \subset (N(f))_i$ . On application of extension procedures, we obtain

$$\varepsilon \Pi_{i=n-2}^n \mu_{i=N(f)_i}^*(E_{k_i}) \leq \int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_{i=N(f)_i}^*(E_{k_i}), Z' > \Rightarrow \varepsilon \Pi_{i=n-2}^n \mu_i^*(N(f)_i \cap E_{k_i}) \leq \int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(E_{k_i}), Z' >.$$

By hypothesis,  $\int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(N(f))_i, Z' > < M$  and  $\Pi_{i=n-2}^n E_{k_i} \ominus \Pi_{i=n-2}^n(N(f))_i$ . Therefore,

$$\varepsilon \chi_{\Pi_{i=n-2}^n \mu_i^*(E_{k_i})} \leq \int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(E_{k_i}), Z' > < M \Rightarrow \Pi_{i=n-2}^n \mu_i^*(E_{k_i}) \leq M/\varepsilon. \text{ Since}$$

$$\Pi_{i=n-2}^n E_i = \cup_{k=1}^\infty \Pi_{i=n-2}^n E_{k_i}, \text{ we have } \Pi_{i=n-2}^n \mu_i^*(E_i) \leq M/\varepsilon \Rightarrow \Pi_{i=n-2}^n E_i \ominus \Pi_{i=n-2}^n(N(f))_i.$$

**Proposition 5** Let  $\Pi_{i=n-2}^n E_i$  be measurably covered by  $\Pi_{i=n-2}^n F_i$ . Suppose  $\Pi_{i=n-2}^n G_i$  is a measurable set such that  $\Pi_{i=n-2}^n G_i \subset \Pi_{i=n-2}^n(F_i - E_i)$ . If  $(a_i, b_i, c_i) : f_n(a_i, b_i, c_i) \geq \varepsilon$  is a finite partition of  $G_i$  for each  $i = n-2, n-1, n$  and  $n \in \aleph$  such that  $f_n \downarrow 0$ , then  $\Pi_{i=n-2}^n((a_i, b_i, c_i) : f_n(a_i, b_i, c_i) \geq \varepsilon)$  is an empty set.

**Proof.** Let  $\Pi_{i=n-2}^n G_i = N(f_1)$ ,  $\Pi_{i=n-2}^n E_i = \Pi_{i=n-2}^n((a_i, b_i, c_i) : f_n(a_i, b_i, c_i) \geq \varepsilon)$  and

$$M = \max(f_1(a_i, b_i, c_i)) \Rightarrow \chi_{\Pi_{i=n-2}^n G_i} f_1 \leq f_1. \text{ Since } f_n \downarrow 0, \text{ it follows that } f_n \leq M \chi_{\Pi_{i=n-2}^n G_i} \text{ for all } n. \text{ So}$$

$$N(f_n) \subset N(f_1) = \Pi_{i=n-2}^n G_i \Rightarrow \chi_{\Pi_{i=n-2}^n G_i} f_n = f_n. \text{ (see [2, p. 5]) Since } \Pi_{i=n-2}^n E_i \text{ is a finite partition of } \Pi_{i=n-2}^n G_i, \text{ it}$$

follows that  $\Pi_{i=n-2}^n G_i = \Pi_{i=n-2}^n((G_i - E_i) \cup E_i)$ . Following disjoint partition of  $\Pi_{i=n-2}^n G_i$  (see [4, p. 6]) and on application of measurable cover estimate technique, we obtain

$$\int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(G_i), Z' > = \int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(G_i - E_i), Z' > + \int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(E_i), Z' >.$$

Since  $f_n(a_i, b_i, c_i) < \varepsilon$  on  $\Pi_{i=n-2}^n(G_i - E_i)$ , it follows that  $\int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(G_i - E_i), Z' > \leq \varepsilon$

$$\Pi_{i=n-2}^n \mu_i^*(G_i - E_i) \leq \varepsilon \Pi_{i=n-2}^n \mu_i^*(G_i). \text{ Replacing } \Pi_{i=n-2}^n G_i \text{ with } \Pi_{i=n-2}^n E_i \text{ in the inequality } f_n \leq M \chi_{\Pi_{i=n-2}^n G_i}, \text{ we}$$

obtain  $\chi_{\Pi_{i=n-2}^n E_i} f_n \leq M \chi_{\Pi_{i=n-2}^n E_i}$ . Therefore,

$$\int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(G_i), Z' > \leq \varepsilon \Pi_{i=n-2}^n \mu_i^*(G_i) + M \Pi_{i=n-2}^n \mu_i^*(E_i). \text{ Since } \Pi_{i=n-2}^n G_i \subset \Pi_{i=n-2}^n(F_i - E_i)$$

and  $\Pi_{i=n-2}^n E_i \ominus \Pi_{i=n-2}^n F_i$  (by hypothesis), it follows that  $\Pi_{i=n-2}^n \mu_i^*(G_i) = 0$ . Since  $f_n \downarrow 0$  for each  $n$ , it follows that

$$\int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(G_i), Z' > = 0. \text{ Therefore, } \Pi_{i=n-2}^n E_i = \Pi_{i=n-2}^n((a_i, b_i, c_i) : f_n(a_i, b_i, c_i) \geq \varepsilon) = \emptyset.$$

**Theorem 1** Let  $f_n$  be an integrable function with respect to  $(\Pi_{i=n-2}^n)_\Phi < \mu_i^*, Z' >$ . For each  $\varepsilon > 0$ , the set

$$\Pi_{i=n-2}^n A_i \text{ is measurably covered by } \Pi_{i=n-2}^n E_i = N(f_n) \text{ if } \int \int \int f_n \delta(\Pi_{i=n-2}^n)_\Phi < \mu_i^*(E_i - A_i), Z' > < \varepsilon.$$





**Proof.** Let  $\Pi_{i=n-2}^n E_i = ((a, b, c) : f_n(a, b, c) \neq 0)$ . Then,  $\chi_{\Pi_{i=n-2}^n E_i} f_n \leq f_n$  where  $\mu_i^*(E_i) < \infty$  for  $i = n-2, n-1, n$ . Let  $P_o = \Pi_{i=n-2}^n A_{k_i}$  be a countable partition of  $\Pi_{i=n-2}^n A_i$  such that for every countable partition  $P = \Pi_{i=n-2}^n E_{k_i}$  of  $\Pi_{i=n-2}^n E_i$  in  $\Pi_{i=n-2}^n G(R_i)$  we have  $P_o \Theta P$ . Since  $\Pi_{i=n-2}^n A_i = \cup_{k=1}^{\infty} \Pi_{i=n-2}^n A_{k_i}$  and  $\Pi_{i=n-2}^n E_i = \cup_{k=1}^{\infty} \Pi_{i=n-2}^n E_{k_i}$ , applying measurable cover estimate technique, we obtain  $\sum_{k=1}^{\infty} \iiint f_n \delta(\Pi_{i=n-2}^n)_{\Phi} < \mu_i^*(E_{k_i} - A_{k_i}), Z' > < \varepsilon/3$ . Consider the the partition  $P_{m'} = (\Pi_{i=n-2}^n C_{k_i} : k = 1, 2, \dots)$  of  $P - P_o = \Pi_{i=n-2}^n (E_{k_i} - A_{k_i})$  such that  $\Pi_{i=n-2}^n (E_{k_i} - A_{k_i}) = \cup_{k=1}^{\infty} \Pi_{i=n-2}^n C_{k_i}$  and  $(P - P_o) \Theta P_{m'} \iiint f_n \delta(\Pi_{i=n-2}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} C_{k_i} - (E_{k_i} - A_{k_i})), Z' > < \varepsilon/3$ . There is a sequence  $(P_{k_i})_{k \in \mathbb{N}} = \Pi_{i=n-2}^n B_{k_i}$  of a countable partitions of  $P_{m'}$  in  $\Pi_{i=1}^3 G(R_i)$  such that  $\Pi_{i=n-2}^n C_{k_i} = \cup_{k=1}^{\infty} \Pi_{i=n-2}^n B_{k_i}$  and  $P_{m'} \Theta P_{k_i} \Rightarrow \iiint f_n \delta(\Pi_{i=n-2}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} B_{k_i} - C_{k_i}), Z' > < \varepsilon/3$ . Therefore,  $\iiint f_n \delta(\Pi_{i=n-2}^n)^* < \mu_i^*(E_i - A_i), Z' > \leq \sum_{k=1}^{\infty} \iiint f_n \delta(\Pi_{i=n-2}^n)^* < \mu_i^*(E_{k_i} - A_{k_i}), Z' > + \iiint f_n \delta(\Pi_{i=n-2}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} C_{k_i} - (E_{k_i} - A_{k_i})), Z' > + \iiint f_n \delta(\Pi_{i=n-2}^n)_{\Phi} < \mu_i^*(\cup_{k=1}^{\infty} B_{k_i} - C_{k_i}), Z' > < \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon \Rightarrow \iiint f_n \delta(\Pi_{i=n-2}^n)_{\Phi} < \mu_i^*(E_i - A_i), Z' > < \varepsilon$ .

## REFERENCES

1. Bruckner, B. and Thomson, S. (1997), *Real Analysis* (1<sup>st</sup> Edition), PrenticeHall, 1-713.
2. Dorlas T.C. (2010), *Reminder notes for the course on measures on topological spaces*, Dublin Institute for Advanced Studies, Dublin, Ireland, 10, 1-22.
3. Elias, M. and Rami, S. (2005), *Real Analysis* (3<sup>rd</sup> Edition), Princeton University Press, 1-48.
4. Jimenez F. E. and Sanchez P. E.A. (2012), *Lattice copies of  $\ell^2$  in  $L'$  of a Vector Measure and Strongly Orthogonal Sequences*, Journal of Function spaces and applications, Hindawi Publishing Corporation, 1-15.
5. Sanchez P. E.A. (2003), *Vector measure orthonormal functions and best approximation for the 4-norm*, Arch. Math. 80, 177-190.
6. Yaogan Mensah (2013), *Facts about the Fourier-Stieltjes transform of vector measures on Compact Groups*, International Journal of Analysis and Applications, 19-25.