



## Relationship between Path and Series Representations for the Three Basic Univalent G-functions

Amir Pishkoo<sup>1,2</sup>, Maslina Darus<sup>2</sup>

<sup>1</sup>Nuclear Science Research School (NSTRI)

P.O. Box 14395-836, Tehran, Iran

apishkoo@gmail.com (corresponding author)

<sup>2</sup>School of Mathematical Sciences, Faculty of Science and Technology  
Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor Darul Ehsan, Malaysia

maslina@ukm.my

### ABSTRACT

In this paper we demonstrate how series representation for the three basic univalent  $G$ -functions, namely  $G_{0,2}^{1,0}$ ,  $G_{1,2}^{1,1}$ , and  $G_{1,1}^{1,1}$  can be obtained from their Mellin-Barnes path integral representations. In two special cases, the images of third basic univalent  $G$ -function  $G_{1,1}^{1,1}$  are derived by the Biernacki and Libera operators.

**Keywords:** Meijer's  $G$ -function; Univalent function; Univalent  $G$ -function; Biernacki operator; Libera operator.



## Council for Innovative Research

Peer Review Research Publishing System

**Journal:** Journal of Advances in Mathematics

Vol 6, No 1

[editor@cirworld.com](mailto:editor@cirworld.com)

[www.cirworld.com](http://www.cirworld.com), [member.cirworld.com](http://member.cirworld.com)



## INTRODUCTION

In mathematics, the Meijer's G-function was introduced by Cornelis Simon Meijer(1946) as a very general function intended to include all elementary functions and most of the known special functions, for instance:

- $\sin z = \sqrt{\pi} G_{0,2}^{1,0}(\frac{z^2}{4} |_{\frac{1}{2},0}^-), \quad -\frac{\pi}{2} < \arg z \leq \frac{\pi}{2}$
- $\cos z = \sqrt{\pi} G_{0,2}^{1,0}(\frac{z^2}{4} |_{0,\frac{1}{2}}^-), \quad \forall z$
- $\ln z = G_{2,2}^{1,2}(z - 1 |_{1,0}^{1,1}), \quad \forall z$
- $J_\nu(z) = G_{0,2}^{1,0}(\frac{z^2}{4} |_{\frac{\nu}{2},\frac{-\nu}{2}}^-), \quad -\frac{\pi}{2} < \arg z \leq \frac{\pi}{2}$

A definition of the Meijer's G-function is given by the path integral in the complexplane, called Mellin-Barnes type integral see [1-3]:

$$G_{p,q}^{m,n}(a_1 \dots a_p |_{b_1 \dots b_q} z) = \frac{1}{2\pi i} \int_L \frac{\prod_{j=1}^m \Gamma(b_j - s) \prod_{j=1}^n \Gamma(1 - a_j + s)}{\prod_{j=m+1}^q \Gamma(1 - b_j + s) \prod_{j=n+1}^p \Gamma(a_j - s)} z^s ds. \tag{1.1}$$

For the function

$$G_{p,q}^{m,n}(a_1, \dots, a_p |_{b_1, \dots, b_q} z) \tag{1.2}$$

The integers  $m; n; p; q$  are called orders of the G-function;  $a_p$  and  $b_q$  are called "parameters" and in general, they are complex numbers. The definition holds under the assumptions:  $0 \leq m \leq q$  and  $0 \leq n \leq p$ , where  $m, n, p$ , and  $q$  are integer numbers. In [4] the univalent Meijer's G-functions are classified into three types in the form of the following proposition:

**Proposition 1.1** All of the univalent Meijer's G-functions,  $G_{p,q+1}^{1,p}$ , can be considered as the generalised ( $q$ -tuple) fractional differ-integrals of one of the three simplest univalent G-functions, namely,  $G_{0,2}^{1,0}; G_{1,2}^{1,1}$ ; and  $G_{1,1}^{1,1}$ , depending on whether  $p < q; p = q; p = q + 1$ .

In [5-8] G-functions are directly obtained as the solution in Micro- and Nano-structures, and in physical models such as Diffusion equation, Laplace's equation, and Schrodinger equation, respectively.

The classical Erdélyi-Kober operators  $I_{1,m}^{(\gamma_k),(\delta_k)}$ ;  $m$  transform one univalent Meijer's G-function of the lower rank to another univalent Meijer's G-function of the upper rank as the following lemma:

**Lemma 1.2** Let  $|z| < 1$  ( $|z| < 1$  for  $p = q + 1$ ), then

$$G_{p,q+1}^{1,p}(1-a_1, \dots, 1-a_p |_{0,1-b_1, \dots, 1-b_q} -z) = \begin{cases} I_{1,1}^{a_p-1, b_q-a_p} \{ G_{p-1,q}^{1,p-1}(1-a_1, \dots, 1-a_{p-1} |_{0,1-b_1, \dots, 1-b_{q-1}} -z) \} & \text{if } b_q > a_p \\ D_{1,1}^{b_q-1, a_p-b_q} \{ G_{p-1,q}^{1,p-1}(1-a_1, \dots, 1-a_{p-1} |_{0,1-b_1, \dots, 1-b_{q-1}} -z) \} & \text{if } b_q < a_p \end{cases} \tag{1.3}$$

Under the following conditions:

$$\delta_k > 0, \mu = 1, \gamma_k > -2, \tag{1.4}$$

these operators in the space of analytic functions,  $A$ , maps the class  $A$  onto itself.

$$I_{1,m}^{(\gamma_k),(\delta_k)} f(z) = z \sum_{n=0}^{\infty} a_n \prod_{k=1}^m \frac{\Gamma(\gamma_k + n + 2)}{\Gamma(\gamma_k + \delta_k + n + 2)} z^n \tag{1.5}$$

maps the class  $A$  onto itself [9]. In [10] Kiryakova et al. introduced other form of definitions for well-known operators by using generalised fractional calculus. For instance form of the Biernacki and Libera operators are respectively as follows:



$$Bf(z) = \int_0^z \frac{f(\xi)}{\xi} d\xi = \int_0^1 \frac{f(z\sigma)}{\sigma} d\sigma = I_1^{-1,1} f(z), \quad (1.6)$$

and

$$Lf(z) = \frac{2}{z} \int_0^z f(\xi) d\xi = 2 \int_0^1 f(z\sigma) d\sigma = 2I_1^{0,1} f(z). \quad (1.7)$$

## 2 MAIN RESULTS

### 2.1 The First basic univalent G-function

In the study of basic univalent G-functions we first determine the position of poles and zeroes of functions inside the path integral.

$$G_{0,2}^{1,0}[-b_1, b_2 | z] = \frac{1}{2\pi i} \int_L \frac{\Gamma(b_1 - s) z^s}{\Gamma(1 - b_2 + s)} ds. \quad (2.1)$$

Here we see that:

1. Position of poles  $\Gamma(b_1 - s)$ :  $s = b_1 + n$ ;  $n = 0, 1, 2, \dots$
2. Position of zeroes  $\frac{1}{\Gamma(1 - b_2 + s)}$ :  $s = b_2 - 1 - n$ ;  $n = 0, 1, 2, \dots$

We obtain the following

**Theorem 2.3** If  $L$  in (2.1) is a loop beginning and ending at  $+\infty$ , encircling all poles of  $\Gamma(b_1 - s)$  exactly once in the negative direction, then

$$G_{0,2}^{1,0}[-b_1, b_2 | z] = \sum_{n=0}^{\infty} \frac{1}{\Gamma(1 + b_1 - b_2 + n)} \frac{z^{b_1+n}}{n!}. \quad (2.2)$$

**Proof 1** At a simple pole, the residue of function  $f$  is given by

$$\text{Res}(f, c) = \lim_{s \rightarrow c} (s - c) f(s).$$

So the residue is given by  $\frac{(-1)^{n-1}}{n!}$ . Then by putting  $s = n + b_1$  in expression  $\frac{z^s}{\Gamma(1 - b_2 + s)}$  we obtain (2.2).

**Example 2.1** If  $b_1 = 0$ ;  $b_2 = 1/2$  and  $z \rightarrow \frac{z^2}{4}$  in (2.1), then we get


$$\cos z = G_{0,2}^{1,0}\left[-0, \frac{1}{2} \mid \frac{z^2}{4}\right] = \sqrt{\pi} \left(\frac{1}{2\pi i}\right) \int \frac{\Gamma(-s) z^{2s}}{4\Gamma(\frac{1}{2} + s)} ds. \quad (2.3)$$

**Corollary 2.4** Putting  $b_1 = 0$ ;  $b_2 = 1/2$  and  $z \rightarrow \frac{z^2}{4}$  in (2.2) gives

$$\cos z = G_{0,2}^{1,0}\left[-0, \frac{1}{2} \mid \frac{z^2}{4}\right] = \sum_{n=0}^{\infty} \frac{1}{\Gamma(\frac{1}{2} + n)} \frac{z^{2n}}{4^n n!}. \quad (2.4)$$

while  $\Gamma(\frac{1}{2} + n) = \frac{(2n)!}{4^n n!} \sqrt{\pi} = \frac{(2n-1)!!}{2^n} \sqrt{\pi}$  we obtain

$$\cos z = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n}}{(2n)!}. \quad (2.5)$$



**Figure 1:** Poles and zeroes of  $\cos z = G_{0,2}^{1,0} \left[ \begin{matrix} - \\ 0, \frac{1}{2} \end{matrix} \middle| -\frac{z^2}{4} \right]$

It is noted that the product of all the odd integers up to some odd positive integer  $k$  is called the double factorial of  $k$ , and denoted by  $k!!$ . Next,

**Theorem 2.5** Let  $p=1, q=2$  in Lemma 1.2 such that

$$G_{1,3}^{1,1} [a_1 | b_2, b_3] - z = \Gamma_{1,1}^{-a_1, a_1 - b_3} G_{0,2}^{1,0} [0, b_2] - z,$$

then (2.2) implies that

$$G_{1,3}^{1,1} [a_1 | b_2, b_3] - z = \sum_{n=0}^{\infty} \frac{\Gamma(1 - a_1 + n)}{\Gamma(1 - b_2 + n) \Gamma(1 - b_3 + n)} \frac{z^n}{n!}. \quad (2.6)$$

## 2.2 The second basic univalent G-function

$$G_{1,2}^{1,1} [a_1 | b_1, b_2] z = \frac{1}{2\pi i} \int_L \frac{\Gamma(b_1 - s) \Gamma(1 - a_1 + s) z^s}{\Gamma(1 - b_2 + s)} ds. \quad (2.7)$$

Here we see that:

1. Position of poles  $\Gamma(b_1 - s)$ :  $s = b_1 + n$ ;  $n = 0, 1, 2, \dots$
2. Position of poles  $\Gamma(1 - a_1 + s)$ :  $s = a_1 - 1 - n$ ;  $n = 0, 1, 2, \dots$
3. Position of zeroes  $\frac{1}{\Gamma(1 - b_2 + s)}$ :  $s = b_2 - 1 - n$ ;  $n = 0, 1, 2, \dots$

Here we have

**Theorem 2.6** If  $a_1 - b_1 \neq 1, 2, 3, \dots$ , which implies that no pole of  $\Gamma(b_1 - s)$  coincides with any pole of  $\Gamma(1 - a_1 + s)$ , then

$$G_{1,2}^{1,1} [a_1 | b_1, b_2] z = \sum_{n=0}^{\infty} \frac{\Gamma(1 - a_1 + b_1 + n)}{\Gamma(1 + b_1 - b_2 + n)} \frac{z^{b_1 + n}}{n!}. \quad (2.8)$$


**Proof 2.** At a simple pole, the residue of function  $f$  is given by

$$\text{Res}(f, c) = \lim_{s \rightarrow c} (s - c)f(s).$$

L in (2.7) is a loop beginning and ending at  $+\infty$ , encircling all poles of  $\Gamma(b_1 - s)$  exactly once in the negative direction, but not encircling any pole of  $\Gamma(1 - a_1 + s)$ . So the residue is given by  $\frac{(-1)^{n-1}}{n!}$ . Then by putting  $s = n + 1$  in  $\frac{\Gamma(1 - a_1 + s)z^s}{\Gamma(1 - b_2 + s)}$  we obtain (2.8).

**Example 2.2** If  $a_1 = 1 + i$ ,  $b_1 = 0$  and  $b_2 = -i + 1/2$  in (2.7), then we have

$$G_{1,2}^{1,1} \left[ \begin{matrix} 1+i \\ 0, i+\frac{1}{2} \end{matrix} \middle| z \right] = \frac{1}{2\pi i} \int_L \frac{\Gamma(-s)\Gamma(-i+s)}{\Gamma(\frac{1}{2}+i+s)} z^s ds.$$



**Figure 2:** Poles and zeroes of  $G_{1,2}^{1,1} \left[ \begin{matrix} 1+i \\ 0, i+\frac{1}{2} \end{matrix} \middle| z \right]$

If equal parameters appear among the  $a_p$  and  $b_q$  determining the factors in the numerator and the denominator of the integrand, the fraction can be simplified, and the order of the function thereby be reduced. If  $a_1 = b_2$  then

$$G_{1,2}^{1,1} \left[ \begin{matrix} a_1 \\ b_1, b_2 \end{matrix} \middle| z \right] = G_{0,1}^{1,0} \left[ \begin{matrix} - \\ b_1 \end{matrix} \middle| z \right] = \frac{1}{2\pi i} \int_L \Gamma(b_1 - s) z^s ds. \quad (2.9)$$

Here we see that:

1. Position of poles  $\Gamma(b_1 - s)$ :  $s = b_1 + n$ ;  $n = 0, 1, 2, \dots$
2. Position of zeroes: there are no zeroes.

**Example 2.3** If we put  $b_1 = 0$  in (2.9); then we get exponential function

$$e^{-z} = G_{0,1}^{1,0} [0 | z] = \frac{1}{2\pi i} \int_L \Gamma(-s) z^s ds. \quad (2.10)$$

**Corollary 2.7** Putting  $a_1 = b_2$  and  $b_1 = 0$  in (2.8) verifies (2.10)

$$e^{-z} = G_{0,1}^{1,0} [0 | z] = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Next an interesting result as follows:

**Theorem 2.8** Let  $p = 2$ ,  $q = 2$  in Lemma 1.2 such that

$$G_{2,3}^{1,2} \left[ \begin{matrix} a_1, a_2 \\ 0, b_2, b_3 \end{matrix} \middle| -z \right] = I_{1,1}^{-a_2, a_2 - b_3} G_{1,2}^{1,1} \left[ \begin{matrix} a_1 \\ 0, b_2 \end{matrix} \middle| -z \right],$$



then (2.8) implies that

$$G_{2,3}^{1,2} [a_1, a_2 | -z] = \sum_{n=0}^{\infty} \frac{\Gamma(1-a_1+n)\Gamma(1-a_2+n)}{\Gamma(1-b_2+n)\Gamma(1-b_3+n)} \frac{z^n}{n!} \quad (2.11)$$

**Proof 4** Using (1.5) with  $\gamma_k = -a_2$ ;  $\delta_k = a_2 - b_3$ ;  $m = 1$  and  $f(z) = G_{1,2}^{1,1} [a_1 | z]$  given by (2.8) yields series representation for  $G_{2,3}^{1,2} [a_1, a_2 | -z]$ .

### 2.3 The third basic univalent G-function

We begin with the definition of third basic univalent G-function,  $G_{1,1}^{1,1}$ , as follows:

$$G_{1,1}^{1,1} [a_1 | -z] = \frac{1}{2\pi i} \int_L \Gamma(b_1 - s)\Gamma(1 - a_1 + s)(-z)^s ds. \quad (2.12)$$

Here we see that:

1. Position of poles  $\Gamma(b_1 - s)$  :  $s = b_1 + n$ ;  $n = 0, 1, 2, \dots$  .
2. Position of poles  $\Gamma(1 - a_1 + s)$  :  $s = a_1 - 1 - n$ ;  $n = 0, 1, 2, \dots$  .
3. Position of zeroes: there are no zeroes.

**Theorem 2.9** Let  $a_1 - b_1 \neq 1, 2, 3, \dots$ , which implies that no pole of  $\Gamma(b_1 - s)$  coincides with any pole of  $\Gamma(1 - a_1 + s)$ , then

$$G_{1,1}^{1,1} [a_1 | -z] = \sum_{n=0}^{\infty} \frac{\Gamma(1 - a_1 + b_1 + n)}{n!} z^{b_1 + n} \quad (2.13)$$


**Proof 5** At a simple pole, the residue of function  $f$  is given by

$$Res(f, c) = \lim_{s \rightarrow c} (s - c)f(s).$$

$L$  in (2.12) is a loop beginning and ending at  $+\infty$ , encircling all poles of  $\Gamma(b_1 - s)$  exactly once in the negative direction, but not encircling any pole of  $\Gamma(1 - a_1 + s)$ . So the residue is given with  $\frac{(-1)^{n-1}}{n!}$ . Then by putting  $s = n + 1$  in  $\Gamma(1 - a_1 + s)(-z)^s$  we obtain (2.13).

**Example 2.4** If we put  $a_1 = 0$  and  $b_1 = 1$  then the Koebe function can be obtained

$$K(z) = \frac{z}{(1-z)^2} = G_{1,1}^{1,1} [0 | -z] = \frac{1}{2\pi i} \int_L \Gamma(1-s)\Gamma(1+s)z^s ds. \quad (2.14)$$



**Figure 3:** Poles of Koebe function  $G_{1,1}^{1,1}[0|-z]$

**Corollary 2.10** Putting  $a_1 = 0$  and  $b_1 = 1$  in (2.13) verifies (2.14)

$$G_{1,1}^{1,1}[0|-z] = \sum_{n=0}^{\infty} \frac{\Gamma(2+n)}{n!} z^{n+1} = \sum_{n=0}^{\infty} (n+1)z^{n+1} = z + 2z^2 + 3z^3 + \dots$$

**Theorem 2.11** Let  $p=2$ ;  $q=1$  in Lemma 1.2 such that

$$G_{2,2}^{1,2}[a_1, a_2 | -z] = I_{1,1}^{-a_2, a_2 - b_2} G_{1,1}^{1,1}[a_1 | -z],$$

then (2.13) implies that

$$G_{2,2}^{1,2}[a_1, a_2 | -z] = \sum_{n=0}^{\infty} \frac{\Gamma(1-a_1+n)\Gamma(1-a_2+n)}{\Gamma(1-b_2+n)} \frac{z^n}{n!} \quad (2.15)$$

**Proof 6** Using (1.5) with  $\gamma_k = -a_1$ ;  $\delta_k = a_1 - b_2$ ;  $m = 1$  and  $f(z) = G_{1,1}^{1,1}[a_1 | -z]$  given by (2.13) yields series representation for  $G_{2,2}^{1,2}[a_1, a_2 | -z]$ .

**Corollary 2.12** Putting  $a_2 = 1$  and  $b_2 = 0$  in (2.15), and using of (1.6) gives image of  $G_{1,1}^{1,1}$  by Biernacki operator

$$G_{2,2}^{1,2}[a_1, 1 | -z] = I_{1,1}^{-1,1} G_{1,1}^{1,1}[a_1 | -z] = \sum_{n=0}^{\infty} \Gamma(1-a_1+n) \frac{z^n}{(n+1)!}.$$

**Corollary 2.13** Putting  $a_2 = 0$  and  $b_2 = 0$  in (2.15), and using of (1.7) gives image of  $G_{1,1}^{1,1}$  by Libera operator

$$G_{2,2}^{1,2}[a_1, 0 | -z] = I_{1,1}^{0,1} G_{1,1}^{1,1}[a_1 | -z] = \sum_{n=0}^{\infty} \Gamma(1-a_1+n)(n+1) \frac{z^n}{(n+2)!}.$$

## ACKNOWLEDGMENTS

This work was supported by MOHE with the grant number:ERGS/1/2013/STG06/UKM/01/2.

## REFERENCES

- [1] Klimyik, A. 2001. Meijer G-function, Springer, Berlin, 2001.
- [2] Askey, R. 2010. Meijer G-function. Cambridge: Cambridge University Press.



- [3] Erdélyi, A., Magnus, W., Oberhettinger, F. & Tricomi, F. 1953. Higher transcendental functions. New York: McGraw-Hill.
- [4] Pishkoo, A. & Darus, M. 2012. Fractional differintegral transformations of univalent Meijers G-functions. Journal of Inequalities and Applications (2012): Article 36.
- [5] Pishkoo, A. & Darus, M. 2013. Some Applications of Meijer G-functions as Solutions of Differential Equations in Physical Models Journal of Mathematical Physics, Analysis, Geometry 3 (9): 379-391.
- [6] Pishkoo, A. & Darus, M. 2013. Meijers G-functions (MGFs) in Micro- and Nano-structures. Journal of Computational and Theoretical Nanoscience 10 (10): 2478-2483.
- [7] Pishkoo, A. & Darus, M. 2012. Meijers G-functions as the solution of Schrodinger equation. Proceeding of the 43rd Iranian Mathematics Conference. University of Tabriz, Tabriz, Iran.
- [8] Pishkoo, A. & Darus, M. 2013. G-function solutions for Diffusion and Laplace's equations, Journal of Advances in Mathematics, 4(1): 359-365.
- [9] Kiryakova, V. 1994. Generalized fractional calculus and applications. , Vol. 301. Chapman & Hall/CRC.
- [10] Kiryakova, V. , Saigo, M. & Srivastava, H., 1998. Some criteria for univalence of analytic functions involving generalized fractional calculus operators, Fract. Calc. Appl. Anal. 1 (1):79--104.

