



Existence And Uniqueness Of Solution Of Inhomogeneous Semilinear Evolution Equation With Nonlocal Condition

¹H. L. Tidke, ²R. T. More

¹Department of Mathematics,
North Maharashtra University, Jalgaon-425 001, India
tharibhau@gmail.com

² Department of Mathematics,
Arts, Commerce and Science College, Bodwad, Jalgaon-425 310, India
rupeshmore9@yahoo.com


ABSTRACT

In this paper, we study the existence and uniqueness of solution of inhomogeneous semilinear evolution equation with nonlocal condition in cone metric space. The result is obtained by using the some extensions of Banach's contraction principle in complete cone metric space.

Keywords:

Inhomogeneous semilinear evolution equation; Cone metric space; Fixed point theorem; Nonlocal condition.

Mathematics Subject Classification: 34L30, 58D25, 65J08, 47J35, 74H10, 34B10.



Council for Innovative Research

Peer Review Research Publishing System

Journal: Journal of Advances in Mathematics

Vol 6, No.3

editor@cirworld.com

www.cirworld.com, member.cirworld.com



1. INTRODUCTION

The purpose of this paper is study the existence and uniqueness of solution of inhomogeneous semilinear evolution equation with nonlocal condition in cone metric space of the form:

$$x'(t) = A(t)x(t) + f(t, x(t)), \quad t \in J = [0, b] \quad (1)$$

$$x(0) + g(x) = x_0, \quad (2)$$

where $A(t)$ is a bounded linear operator on a Banach space X with domain $D(A(t))$, the unknown $x(\cdot)$ takes values in the Banach space X ; $f: J \times X \rightarrow X$, $g: C(J, X) \rightarrow X$ are appropriate continuous functions and x_0 is given element of X .

Many authors have been studied the problems of existence, uniqueness, continuation and other properties of solutions of these type or special forms of the equations (1)–(2) are studied by different techniques, for example, see [3, 4, 5, 8, 10, 13] and the references given therein.

The objective of the present paper is to study the existence and uniqueness of solution of the evolution equation (1.1)–(1.2) under the conditions in respect of the cone metric space and fixed point theory. Hence we extend and improve some results reported in [2, 8, 10, 11, 13].

The paper is organized as follows: Section 2, we discuss the preliminaries. Section 3, we dealt with study of existence and uniqueness of solution of inhomogeneous evolution equation with nonlocal condition in cone metric space. Finally in Section 4, we give example to illustrate the application of our result.

2. Preliminaries

Let us recall the concepts of the cone metric space and we refer the reader to [1, 6, 7, 9, 12] for the more details.

Let E be a real Banach space and P is a subset of E . Then P is called a cone if and only if,

1. P is closed, nonempty and $P \neq \{0\}$;
2. $a, b \in \mathbb{R}$, $a, b \geq 0$, $x, y \in P \Rightarrow ax + by \in P$;
3. $x \in P$ and $-x \in P \Rightarrow x = 0$.

For a given cone $P \subset E$, we define a partial ordering relation \leq with respect to P by $x \leq y$ if and only if $y - x \in P$. We shall write $x < y$ to indicate that $x \leq y$ but $x \neq y$, while $x \ll y$ will stand for $y - x \in \text{int}P$, where $\text{int}P$ denotes the interior of P .

The cone P is called normal if there is a number $K > 0$ such that $0 \leq x \leq y$ implies $\|x\| \leq K\|y\|$, for every $x, y \in E$. The least positive number satisfying above is called the normal constant of P .

In the following we always suppose E is a real Banach space, P is a cone in E with $\text{int}P \neq \emptyset$, and \leq is partial ordering with respect to P .

Definition 2.1 Let X be a nonempty set. Suppose that the mapping $d: X \times X \rightarrow E$ satisfies:

1. $0 \leq d(x, y)$ for all $x, y \in X$ and $d(x, y) = 0$ if and only if $x = y$;
2. $d(x, y) = d(y, x)$, for all $x, y \in X$;
3. $d(x, y) \leq d(x, z) + d(z, y)$, for all $x, y, z \in X$.

Then d is called a cone metric on X and (X, d) is called a cone metric space. The concept of cone metric space is more general than that of metric space.

The following example is a cone metric space, see [11].

Example 2.2 Let $E = \mathbb{R}^2$, $P = \{(x, y) \in E : x, y \geq 0\}$, $X = \mathbb{R}$, and $d: X \times X \rightarrow E$ such that $d(x, y) = (|x - y|, \alpha|x - y|)$, where $\alpha \geq 0$ is a constant. Then (X, d) is a cone metric space.



Definition 2.3 Let X be an ordered space. A function $\Phi: X \rightarrow X$ is said to be a comparison function if for every $x, y \in X$, $x \leq y$, implies that $\Phi(x) \leq \Phi(y)$, $\Phi(x) \leq x$ and $\lim_{n \rightarrow \infty} \|\Phi^n(x)\| = 0$, for every $x \in X$.

Example 2.4 Let $E = \mathbb{R}^2$, $P = \{(x, y) \in E : x, y \geq 0\}$. It is easy to check that $\Phi: E \rightarrow E$, with $\Phi(x, y) = (ax, ay)$, for some $a \in (0, 1)$ is a comparison function. Also if Φ_1, Φ_2 are two comparison functions over \mathbb{R} , then $\Phi(x, y) = (\Phi_1(x), \Phi_2(y))$ is also a comparison function over E .

3. Existence and uniqueness of solution

Let X be a Banach space with norm $\|\cdot\|$. Let $B = C(J, X)$ be the Banach space of all continuous functions from J into X endowed with supremum norm

$$\|x\|_{\infty} = \sup\{\|x(t)\| : t \in J\}.$$

Let $P = \{(x, y) : x, y \geq 0\} \subset E = \mathbb{R}^2$ be a cone and define $d(f, g) = (\|f - g\|_{\infty}, \alpha\|f - g\|_{\infty})$, for every $f, g \in B$. Then it is easily seen that (B, d) is a cone metric space.

Definition 3.1 The function $x \in B$ satisfies the integral equation

$$x(t) = x_0 - g(x) + \int_0^t A(s)f(s, x(s))ds, \quad t \in J \quad (3)$$

is called the solution of the evolution equation (1)–(2).

We need the following lemma for further discussion:

Lemma 3.2 [11] Let (X, d) be a complete cone metric space, where P is a normal cone with normal constant K . Let $f: X \rightarrow X$ be a function such that there exists a comparison function $\Phi: P \rightarrow P$ such that

$$d(f(x), f(y)) \leq \Phi(d(x, y)),$$

for every $x, y \in X$. Then f has a unique fixed point.

We list the following hypotheses for our convenience:

(H₁) $A(t)$ is a bounded linear operator on X for each $t \in J$, the function $t \rightarrow A(t)$ is continuous in the uniform operator topology and hence there exists a constant K such that

$$K = \sup_{t \in J} \|A(t)\|.$$

(H₂) There exist continuous function $p: J \rightarrow \mathbb{R}^+$ and a comparison function $\Phi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

$$(\|f(t, x) - f(t, y)\|, \alpha\|f(t, x) - f(t, y)\|) \leq p(t)\Phi(d(x, y)),$$

and for positive constant G , such that

$$(\|g(x) - g(y)\|, \alpha\|g(x) - g(y)\|) \leq G\Phi(d(x, y)),$$

for every $t \in J$ and $x, y \in X$.

(H₃) $\sup_{t \in J} [G + \int_0^t Kp(s)ds] = 1$.

Theorem 3.3 Assume that hypotheses (H₁)–(H₃) hold. Then the evolution equation (1.1)–(1.2) has a unique solution x on J .

Proof: The operator $F: B \rightarrow B$ is defined by

$$Fx(t) = x_0 - g(x) + \int_0^t A(s)f(s, x(s))ds, \quad t \in J. \quad (4)$$



By using the hypotheses $(H_1) - (H_3)$, we have

$$\begin{aligned}
 (\|Fx(t) - Fy(t)\|, \alpha\|Fx(t) - Fy(t)\|) &\leq (\|g(x) - g(y)\| + \int_0^t A(s)\|f(s, x(s)) - f(s, y(s))\| ds, \\
 &\quad \alpha\|g(x) - g(y)\| + \alpha \int_0^t A(s)\|f(s, x(s)) - f(s, y(s))\| ds) \\
 &\leq (\|g(x) - g(y)\|, \alpha\|g(x) - g(y)\|) \\
 &\quad + \int_0^t K(\|f(s, x(s)) - f(s, y(s))\|, \alpha\|f(s, x(s)) - f(s, y(s))\|) ds \\
 &\leq G\Phi(\|x - y\|, \alpha\|x - y\|) + \int_0^t Kp(s)\Phi(\|x(s) - y(s)\|, \alpha\|x(s) - y(s)\|) ds \\
 &\leq G\Phi(\|x - y\|_b, \alpha\|x - y\|_b) + \Phi(\|x - y\|_b, \alpha\|x - y\|_b) \int_0^t Kp(s) ds \\
 &\leq G\Phi(d(x, y)) + \Phi(d(x, y)) \int_0^t Kp(s) ds \\
 &\leq \Phi(d(x, y)) [G + \int_0^t Kp(s) ds] \\
 &\leq \Phi(d(x, y)), \tag{5}
 \end{aligned}$$

for every $x, y \in B$. This implies that $d(Fx, Fy) \leq \Phi(d(x, y))$, for every $x, y \in B$. Now an application of Lemma 3.2, the operator has a unique point in B . This means that the equation (1)–(2) has unique solution. This completes the proof of the Theorem 3.3.

4. Application

In this section, we give an example to illustrate the usefulness of our result discussed in previous section. Let us consider the following evolution equation:

$$\frac{dx}{dt} = \frac{140}{8} e^{-t} x(t) + \frac{te^{-t} x(t)}{(9 + e^t)(1 + x(t))}, \quad t \in J = [0, 1], \quad x \in X, \tag{6}$$

$$x(0) + \frac{x}{8 + x} = x_0, \tag{7}$$

Therefore, we have

$$\begin{aligned}
 A(t) &= \frac{140}{8} e^{-t}, \\
 f(t, x(t)) &= \frac{te^{-t} x(t)}{(9 + e^t)(1 + x(t))}, \quad (t, x) \in J \times X \\
 g(x) &= \frac{x}{8 + x}, \quad x \in X.
 \end{aligned}$$

Now for $x, y \in C(J, X)$ and $t \in J$, we have

$$\begin{aligned}
 (\|Fx(t) - Fy(t)\|, \alpha\|Fx(t) - Fy(t)\|) &= \frac{te^{-t}}{9 + e^t} (\|\frac{x(t)}{1 + x(t)} - \frac{y(t)}{1 + y(t)}\|, \alpha\|\frac{x(t)}{1 + x(t)} - \frac{y(t)}{1 + y(t)}\|) \\
 &= \frac{te^{-t}}{9 + e^t} (\|\frac{x(t) - y(t)}{(1 + x(t))(1 + y(t))}\|, \alpha\|\frac{x(t) - y(t)}{(1 + x(t))(1 + y(t))}\|)
 \end{aligned}$$



$$\begin{aligned}
&\leq \frac{te^{-t}}{9+e^t} (\|x(t) - y(t)\|, \alpha \|x(t) - y(t)\|) \\
&\leq \frac{te^{-t}}{9+e^t} (\|x - y\|_\infty, \alpha \|x - y\|_\infty) \\
&\leq \frac{te^{-t}}{9+e^t} d(x, y) \\
&\leq \frac{t}{10} \Phi(d(x, y)), \tag{8}
\end{aligned}$$

where $p(t) = \frac{t}{10}$, which is continuous function of J into \mathbb{R}^+ and a comparison function $\Phi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $\Phi(d(x, y)) = d(x, y)$.

Similarly, we can have

$$\begin{aligned}
(\|g(x) - g(y)\|, \alpha \|g(x) - g(y)\|) &\leq 8 \left(\frac{\|x - y\|}{(8 + \|x\|)(8 + \|y\|)}, \alpha \frac{\|x - y\|}{(8 + \|x\|)(8 + \|y\|)} \right) \\
&\leq \frac{8}{64} (\|x - y\|, \alpha \|x - y\|) \\
&\leq \frac{1}{8} (\|x - y\|_\infty, \alpha \|x - y\|_\infty) \\
&\leq \frac{1}{8} \Phi(d(x, y)), \tag{9}
\end{aligned}$$

where $G = \frac{1}{8}$, and the comparison function Φ defined as above. Hence the condition (H_1) holds with $K = \frac{140}{8}$.

Moreover,

$$\begin{aligned}
\sup_{t \in J} [G + K \int_0^t p(s) ds] &= \sup_{t \in J} \left[\frac{1}{8} + \frac{140}{8} \int_0^t \frac{s}{10} ds \right] \\
&= \sup_{t \in J} \left[\frac{1}{8} + \frac{140}{8} \frac{t^2}{20} \right] \\
&= \sup_{t \in J} \left[\frac{1}{8} + \frac{7}{8} t^2 \right] \\
&= \left[\frac{1}{8} + \frac{7}{8} \right] = 1. \tag{10}
\end{aligned}$$

Since all the conditions of Theorem 3.3 are satisfied, the problem (6)–(7) has a unique solution x on J .

REFERENCES

- [1] Abbas, M. and Jungck, G. 2008. Common fixed point results for noncommuting mappings without continuity in cone metric spaces, *Journal of Mathematical Analysis and Applications*, Vol. 341, No.1, 416-420.
- [2] Banas, J. 2006. Solutions of a functional integral equation in BC, *International Mathematical Forum*, 1 (2006), No. 24, 1181-1194.
- [3] Burton, T. A. 1983. *Volterra Integral and Differential Equations*, Academic Press, New York, (1983).



- [4] Coddington, E. A. 2003. An Introduction to Ordinary Differential Equations, Prentice-Hall of India, Private Limited, New Delhi, (2003).
- [5] Deo, S. G., Lakshmikantham, V. and Raghavendra, V. 2003. Text Book of Ordinary Differential Equations, Tata McGraw-Hill Publishing Company Limited, New Delhi, Edition, (2003).
- [6] Huang, L. G. and Zhang, X. 2007. Cone metric spaces and fixed point theorems of contractive mappings, Journal of Mathematical Analysis and Applications, Vol. 332, (2007), No.2, 1468-1476.
- [7] Ilic, D. and Rakocevic, V. 2008. Common fixed points for maps on cone metric space, Journal of Mathematical Analysis and Applications, Vol. 341, (2008), No.2, 876-882.
- [8] Karoui, A. 2005. On the existence of continuous solutions of nonlinear integral equations, Applied Mathematics Letters, 18(2005), 299-305.
- [9] Kwong, M. K, 2008. On Krasnoselskii's cone fixed point theorems, Fixed Point Theory and Applications, Volume 2008, Article ID 164537, 18 pages.
- [10] Pazy, A. 1983. Semigroups of Linear Operators and applications to Partial Differential Equations, Springer-Verlag, New York, (1983).
- [11] Raja, P. and Vaezpour, S. M. 2008. Some extensions of Banach's contraction principle in complete cone metric spaces, Fixed Point Theory and Applications, Volume 2008, Article ID 768294, 11pages.
- [12] Rezapour, Sh. and Hamlbarani, R. 2008. Some notes on the paper "Cone metric spaces and fixed point theorems of contractive mappings", Journal of Mathematical Analysis and Applications, Vol. 345, (2008), No.2, 719-724.
- [13] Tidke, H. L. and Dhakne, M. B. 2010. Existence and uniqueness of solution of differential equation of second order in cone metric spaces, Fasciculi Mathematici, Nr. 45, (2010), 121-131.

